

## Efficiency of propolis extract against mitochondrial stress induced by antineoplastic agents (doxorubicin and vinblastin) in rats

Lamia Benguedouar, Hanane Nadia Boussenane, Wided Kebsa,  
Mohamed Alyane, Hassiba Rouibah & Mesbah Lahouel\*

Laboratoire de Pharmacologie et Phytochimie, Faculté des Sciences,  
Université de Jijel. 18000 Jijel, Algérie

Received 09 April 2007 ; revised 06 November 2007

To assess the oxidative stress and mitochondrial dysfunction associated with disease, toxic process and aging, *in vivo* and *in vitro* preventive effect of propolis extract against mitochondrial oxidative stress induced by two anticancer drugs (doxorubicin and vinblastin) have been investigated in female wistar rat using liver and heart mitochondria. The results show that doxorubicin and vinblastin altered mitochondrial functions as observed by a decrease in respiratory control value, an activation of swelling and overproduction of superoxide anion. Myocardial tissue from doxorubicin treated rats showed a marked increase in malondialdehyde production, a depletion of reduced glutathione contents and an inhibition of catalase and superoxide dismutase activities. Similar results were also observed in liver tissue. Pretreatment of rats with propolis extract (100 mg/kg/day po) ( $10^{-4}$  M ip) administered 4 days prior to doxorubicin (20 mg/kg) and/or vinblastin (2 mg/kg) injection, substantially reduced the peroxidative damage in myocardium and hepatic tissues and markedly restored the tissues catalase and SOD activities. The results strongly suggest that propolis extract protects heart and liver tissues from oxidative stress by protecting the mitochondria.

**Keywords:** Doxorubicin, Heart, Liver, Lipid peroxidation, Propolis extract, Vinblastin

Doxorubicin and vinblastin are important and effective anticancer drugs widely used for the treatment of various types of cancer but their clinical uses are limited by dose-dependent cardiotoxicity and hepatotoxicity. Propolis extract protects, *in vivo*, the liver and blood tissues against cyclophosphamid and doxorubicin toxicity<sup>1,2</sup> suggesting that polyphenols can help in preventing the oxidative stress induced by chemotherapeutic agents.

Oxidative stress and mitochondrial dysfunction are associated with disease and aging<sup>3</sup>. Oxidative stress results from an over production of reactive oxygen species (ROS), often leading to peroxidation of membrane phospholipids and production of reactive aldehydes. Moreover, inner mitochondria membrane is considered as targets of apoptotic process, owing to alterations of potential membrane<sup>4-7</sup> and to ROS production.

Polyphenolic compounds, including flavonoids, tannins and phenolic acids are widely distributed in

plant and foods of plant origin<sup>8</sup>. The chemical properties of phenolic acids or flavonoids, in terms of availability of the phenolic hydrogens as hydrogen donating radical scavengers, predict their antioxidant properties<sup>9</sup>.

Considerable interest in these compounds extracted from propolis has arisen with the recognition for their antibacterial and antioxidant effects<sup>10-12</sup>. However despite antioxidant and antibacterial data on the beneficial effects of propolis extracts, the molecular mechanisms underlying interactions with isolated mitochondria have not yet been reported. Some of their protective effect can be attributed to direct scavenging properties<sup>13,14</sup>. Mitochondrial target was evoked to explain the protecting effect of polyphenols<sup>15-17</sup>. Some polyphenols with antioxidant and free radical scavenging properties were shown to inhibit also lipid peroxidation (Resveratrol, Propolis, *Ginkgo biloba*)<sup>1,18,19</sup>. This protective effect could be due to a direct action on mitochondrial functions.

With a view to elucidate the mechanism of interaction of polyphenols present in propolis on heart and liver mitochondria, the present study reports mitochondrial functions following chemotherapeutic drugs toxicity in female rats.

\*Correspondent author

Telephone : (+213) 34 47 24 37

Fax : (+213) 34 47 48 96

E-mail : lahoulmesbah@yahoo.fr

## Materials and Methods

**Chemicals**—Sucrose, EGTA, rotenone, malonate, antimycin A, oligomycin, succinate, malate, glutamate, NADH, ADP, Tris base and  $\text{KH}_2\text{PO}_4$ , cyclosporine A, albumin, pyruvate, succinate, nitroblue tetrazolium (NBT), vitamin E, quercetin, 1,1-diphenyl-2-picrylhydrazyl hydrate (DPPH) and dithiobisnitrobenzoic acid (DTNB) were purchased from Sigma Chemical Co. KCl,  $\text{MgCl}_2$  and KCN were obtained from Prolabo. All drugs were solubilized in distilled water in order to obtain a stock solution of  $10^{-3}$  M. N,N,N',N'-tetramethyl-*p*-phenylenediamine and oligomycin were solubilized in dimethylformamide (DMF). All the controls experiments were carried out using the same mixture. The propolis extract solubilized in ethanol was added at different final concentrations, between  $10^{-3}$  M to  $10^{-9}$  M (quercetin equivalent).

**Preparation of propolis extract and flavonoid assessment**—Propolis extract was obtained by using ethanol and methanol. The propolis (beeswax and resins collected by the honeybee from plants, particularly from flowers and leaf buds, collected from the region of Jijel, Algeria), was cut into small pieces; dived in 95 % alcohol for 15 days, filtered and then evaporated to  $80^\circ\text{C}$  using a rotary evaporator (Evaporator E100). The residual was retaken using 70 % methanol and was let for steeping for one night. After evaporation of the solvent, the product extract is called “raw extract or propolis extract”. Flavonoids rate was determined by reactivity with  $\text{AlCl}_3$  method as described by Bankova<sup>20</sup>. The concentrations of propolis extract were given as concentration quercetin equivalent ( $10^{-4}$  M).

**Free radical scavenging assay**—The free radical-scavenging activity of test compounds (propolis extract, vitamin E and quercetin) was evaluated using the stable radical DPPH $\cdot$ . 15  $\mu\text{l}$  of each test compound was added to 1.5 ml (100  $\mu\text{M}$ ) DPPH $\cdot$  in methanol. The absorbancy of the remaining DPPH $\cdot$  was determined at 15 sec interval during 5 min at 515 nm. Blank sample contained the same amount of methanol and DPPH $\cdot$ . The measurements were performed in triplicate. The radical scavenging activity was calculated by the formula:

$$I = [(A_B - A_A)/A_B] 100$$

where: I = DPPH $\cdot$  inhibition, (%);  $A_B$  = absorption of blank sample (at  $t = 0$  min) and  $A_A$  = absorption of tested compounds.

**Animals treatment**—Female Wistar rats (purchased from Pasteur Institute, Algiers, Algeria) weighing 250-280 g were housed in stainless steel cages in room with a controlled temperature ( $22^\circ\text{C}$ ) and light cycle light on at 1200 hr) during the experimental period. They had free access to food (Groupe Avicole, Medea, Algeria) and water. The rats were divided into three subgroups:

Pretreated group ( $n = 24$ ), rats were treated with propolis extract (100 mg/kg/day, po) for 4 days prior to anticancer drug administration. After 4 days, they were injected, ip, with adriamycin (20 mg/kg), vinblastin (2 mg/kg) or adriamycin (20 mg/kg) + vinblastin (2 mg/kg).

Untreated group ( $n = 24$ ), rats were injected, ip, with adriamycin (20 mg/kg), vinblastin (2 mg/kg) or adriamycin (20 mg/kg) + vinblastin (2 mg/kg).

Control group ( $n = 8$ ), rats were injected, ip, with 0.9 % saline solution.

After 24 hr of injections, the rats were sacrificed by decapitation and their livers and hearts were rapidly excised, weighed and placed in ice-bath.

**Isolation of rat liver and heart mitochondria**—Mitochondria were extracted from a homogenate of rat heart and liver by differential centrifugations<sup>21</sup>. Briefly, samples of freshly excised hearts (approximately 0.5 g) and livers (approximately 5 g) were respectively homogenized in ice-cold heart isolation medium (mannitol 225 mM, sucrose 75 mM, Hepes 5 mM, EGTA 1 mM and albumin 1 mg/ml, pH 7.4 at  $37^\circ\text{C}$ ) and liver isolation medium (Tris 20 mM, sucrose 250 mM, EGTA 2 mM and  $\text{MgCl}_2$  5 mM, pH 7.2 at  $4^\circ\text{C}$ ) using a Potter-Elvehjem homogenizer. The homogenates were centrifuged at 2000 g for 10 min at  $4^\circ\text{C}$  to remove cell debris and nuclei and supernatants were collected. Mitochondria were separated from the supernatant by centrifugation at 12000 g for 10 min. The pellet (mitochondria) was washed and resuspended in a respiratory buffer ( $\text{KH}_2\text{PO}_4$  1 mM, sucrose 50 mM, KCl 100 mM, pH 7.4 at  $37^\circ\text{C}$ ) or ( $\text{KH}_2\text{PO}_4$  10 mM, sucrose 200 mM, pH 7.2 at  $25^\circ\text{C}$ ) for measuring heart and liver mitochondrial enzymatic and respiratory activities respectively. Protein concentration of the mitochondrial suspension was assayed by Lowry's method.

**Measurement of mitochondrial respiration**—Oxygen uptake was determined with a Clark-type microelectrode (Hansatech). Each experiment was carried out as follows: aliquot of mitochondria suspension equivalent to 1mg/ml were incubated with

(or without) the tested drug for 1 min at 37°C or 25°C in 1000 µl of the respiratory buffer with or without the inhibitors, then the substrate (pyruvate/malate 6 mM for heart mitochondria) or (succinate 6 mM for liver mitochondria) was added and oxygen consumption was checked (State 2). To initiate state 3 respiratory activity, 200 µM ADP was added to the cuvette. When all ADP was converted to ATP, the state 4 was measured. The following parameters were determined: the respiratory rates calculated as nanomoles of O<sub>2</sub>/min/ mg of mitochondrial protein and respiratory chain ratio (RCR) expressed as the ratio of state 3/state 4 oxygen consumption.

**Mitochondrial superoxide anion (O<sub>2</sub><sup>-</sup>) assay**—Generation of O<sub>2</sub><sup>-</sup> was achieved using rat heart and liver mitochondria. Reaction mixture contained mitochondria (0.2 mg/ml), 1 µM CA and NBT (100 µM) in 1.2 ml of respiration buffer at 37°C and 25°C. The reaction was started by adding malate/pyruvate (6 mM) or succinate (6 mM) for heart and liver respectively in the assay cuvette and the rate of NBT reduction was measured at 560 nm. In absence of mitochondrial substrate (malate/pyruvate) or (succinate), there was no O<sub>2</sub><sup>-</sup> production.

**Mitochondrial swelling measurement**—Mitochondrial swelling was assessed by measuring the change in absorbance of the suspension at 540 nm (A<sub>540</sub>) using a Hitachi model UV-3000 spectrophotometer. Mitochondria (1 mg/ml) were incubated in a total volume of 1.8 ml of heart mitochondria buffer (KH<sub>2</sub>PO<sub>4</sub> 1 mM, Sucrose 50 mM, KCl 100 mM, pH 7.4 at 37°C) in the presence of malate/pyruvate 6 mM at 37°C or in ice-cold isolation liver mitochondria medium (Tris 20 mM, sucrose 250 mM, EGTA 2 mM and MgCl<sub>2</sub> 5 mM, pH 7.2 at 4°C). 100 µM CaCl<sub>2</sub> was added to induce swelling before addition of increasing concentrations of the propolis extract.

**Lipid peroxidation assessment**—The lipid peroxidation in heart and liver mitochondria was assessed using an incubation mixture, containing 800 µl of suspended mitochondria in 0.9 % NaCl, 100 µl of the oxidizing solution Fe<sup>2+</sup>/Fe<sup>3+</sup> (50 µM / 50 µM). Incubation was carried out in a water bath at 37°C for 30 min. The reaction was stopped by the addition of 1 ml of 10 % TCA. The tubes were shaken well and 1.5 ml of thiobarbituric acid (1 % in 0.05 NaOH) reagent was added and the mixture was heated at 100°C for 30 min. Tubes were recooled in ice for 5 min, then centrifuged at 3000 rpm for 15 min and the

colours developed in the supernatant were read at 530 nm. The reaction without Fe<sup>2+</sup>/Fe<sup>3+</sup> was carried out for each mitochondria suspension as the blank. Results were expressed as µM MDA / mg of mitochondrial membrane proteins.

**Dosage of glutathione (GSH)**—Portions (approximately 1 g of liver or 0.5 g of heart freshly excised or frozen) were homogenized in 3 volumes of 5% TCA using Dounce homogenizer. The samples were centrifuged at 2000 rpm for 15 min. The supernatant (50µl) was diluted in 10 ml phosphate buffer (0.1 M, pH 8). Consequently, 20 µl of DTNB 0.01 M was added to 3 ml of the dilution mixture. The measurement was performed at 412 nm against a control prepared in the same conditions using 5% TCA. The concentrations are expressed in mmoles of GSH / g of liver. They are deducted from a range of GSH, which was prepared with the same conditions as dosage did.

**Catalase and superoxide dismutase (SOD) measurement**—Isolated mitochondria prepared as described above were sonicated for 1 min. The sonic extract was centrifuged at 12,000 rpm for 15 min. The supernatant was then assayed by spectrophotometry for catalase activity by the method of Clairborne<sup>22</sup> using catalase (Sigma Chemical Co.) as a standard and the SOD activity by the method of Beauchamp<sup>23</sup> using SOD (Sigma Chemical Co.) as a standard.

**Statistical analysis**—The data are presented as mean±SD of three experiments for each compound and different experimental group. The Student's *t* test and/or the F test (one way analysis of variance [ANOVA]) were used to evaluate the probability of significant differences between groups and compounds. Significance was considered to have occurred at *P* < 0.05.

## Results

Doxorubicin has no effects on liver and vinblastin on heart mitochondria. Therefore, the data are not presented.

**Antioxidant effects of propolis extract**—There are many different antioxidant components in a natural sample, and it is relatively difficult to measure each one separately. Several methods have been employed to determine antioxidant activity of biological samples, and the results are compared with those of reference antioxidant standards.

Free radical-scavenging activity of the propolis extract using DPPH<sup>•</sup> was investigated. The results were compared with those of reference antioxidants.

A higher extract concentration required to scavenge the radicals means a lower antioxidant activity. The decrease of absorbance at 515 nm with the presence of antioxidants indicates the consumption of free radicals in the reaction mixture. The propolis ethanol extract at  $10^{-4}$  M exhibited the highest free radical scavenging activity when compared with the other two polyphenols, vitamin E and quercetin (Fig.1).

*Effect of propolis extract on SO production*—In order to evaluate antioxidant role of the propolis extract in heart and liver mitochondria, SO anions production was measured *in vitro* and *in vivo* on isolated heart and liver mitochondria. Propolis extract reduced the production of anion at concentrations between  $10^{-4}$  -  $10^{-8}$  M *in vitro* either on mitochondria incubated with doxorubicin 1  $\mu$ M or control (Fig.2). *In vivo*, propolis extract (100 mg/kg) given orally, starting 4 days prior to the administration of the anticancer drugs (doxorubicin and vinblastin) also reduced significantly the heart and liver mitochondrial production of SO ( $P < 0.01$ ) (Figs. 3 a and b).

*Effect of propolis extract on lipid peroxidation*—Propolis extract inhibits the lipid peroxidation of heart




Fig. 1—Flavonoids scavenger effects (DPPH<sup>+</sup> Reduction).




Fig. 2—Effect of different concentrations ( $10^{-2}$  -  $10^{-10}$  M) of propolis extract on the heart mitochondrial production of SO *in vitro*.

or liver mitochondrial and cellular membranes (Fig.4). The inhibition of lipid peroxidation reached 90% at  $10^{-6}$  M concentration. *In vitro*, the inhibitory effect of propolis extract on lipid peroxidation of heart and liver mitochondrial membranes was observed at  $10^{-4}$  M showing it to be more than a ten stronger than that of the positive control, quercetin. *In*




Fig. 3—a: Effect of (i) doxorubicin (20 mg/kg) and (ii) the propolis extract on the heart mitochondrial production of SO *in vivo*. b: Effect of (i) doxorubicin (20 mg/kg)+vinblastin (2 mg/kg) treatment and (ii) the propolis extract pre-treatment on the liver mitochondrial production of SO *in vivo*.




Fig. 4—Variations of heart and liver mitochondria MDA levels (nMole/mg protein) expressed as % inhibition of control in the presence of different concentrations (ranged between  $10^{-2}$  and  $10^{-10}$  molar) of propolis extract.

*in vivo*, a significant reduction of MDA concentrations in heart and liver cells of animals pre-treated with 100 mg/kg propolis extract daily for five days prior to doxorubicin (20 mg/kg) or doxorubicin (2 mg/kg) + vinblastin (2 mg/kg) iv injection was observed (Table 1).

*Effect of propolis extract on mitochondrial respiration and swelling*—It was observed that 20




Fig. 5—a: Effects of doxorubicin (20 mg/kg) doxorubicin (20 mg/kg)+vinblastin (2 mg/kg) alone or in association with the propolis extract on the heart and liver mitochondrial respiration function respectively. The values are mean of three experiments for each animal (n = 10). b: Effect of doxorubicin (20 mg/kg), doxorubicin (20 mg/kg)+vinblastin (2 mg/kg) administrations alone or in association with the propolis extract on the heart and liver mitochondrial P/O ratio respectively. The values are mean of three experiments for each animal (n = 10) [i = heart control, ii = doxorubicin (20 mg/kg), iii = propolis+doxorubicin, iv = liver control, v = doxorubicin (20 mg/kg)+vinblastin (2 mg/kg), vi = propolis+doxorubicin+vinblastin].

Table 1—Effect of polyphenols (propolis extract) on the cellular lipid peroxidation induced by doxorubicin (20 mg/kg) and vinblastin (2 mg/kg) (\*\*\*)  $P < 0.001$

[Values are mean  $\pm$  SD]

	MDA (nM/mg heart cell proteins)	MDA (nM/mg liver cell proteins)
Control (n = 8)	11.53 $\pm$ 3.50	31.53 $\pm$ 3.50
Doxorubicin (20 mg/kg) + Vinblastin (2 mg/kg) (n = 5)	19.08 $\pm$ 4.12 *	49.00 $\pm$ 6.71 *
Propolis 100 mg/kg + Doxorubicin 20 mg/kg (n = 8)	10.15 $\pm$ 2.65 *	28.40 $\pm$ 5.50 *

\* $P < 0.001$

mg/kg doxorubicin and 20 mg/kg doxorubicin + 2 mg/kg vinblastin increased the oxygen consumption (state 3) and the inhibition of state 4 occurring a decrease of respiratory chain ratio (RCR =  $V_3 / V_4$ ) and a decrease of P/O also (Figs 5a and b). Propolis extract inhibited the effect of drugs showing a restoration of RCR and P/O *in vitro* and *in vivo* ( $P < 0.001$ ). Additional propolis extract to a mixture of rat heart and liver mitochondria with 25  $\mu$ M calcium, inhibited slightly the rate and amplitude of mitochondrial swelling at  $10^{-5}$  and  $10^{-6}$  M concentrations affected by the doxorubicin and doxorubicin + vinblastin addition to the medium at a concentration of 1  $\mu$ M.

*Effect of propolis extract on GSH reserve*—The effect of chemotherapeutic agents alone or in combination with propolis extract on the tissue levels of GSH is presented in Table 2. An increase of GSH was observed in the group of rats receiving the extract of propolis, compared with controls, while the rats treated by doxorubicin or doxorubicin + vinblastin only showed a decrease of 60 % in GSH levels ( $P < 0.001$ ).

*Effect of propolis extract on enzymes activities; catalase and SOD*—Modification of the mitochondrial enzymes is a signal of anticancer drugs (doxorubicin and vinblastin) oxidative stress. However the propolis extract seems to play an original role to restore the loss of the enzymes activities (catalase and SOD) induced by doxorubicin and vinblastin (Figs 6 and 7). This effect was in relation with the antioxidative effects of polyphenols. Indeed, this result show that polyphenols are able to deactivate quickly the ROS produced by mitochondria.

## Discussion

Oxidative stress and mitochondrial dysfunction result from overproduction by complexes of the respiratory chain of ROS, often leading to peroxidation of membrane phospholipids and production of reactive aldehydes<sup>3,15,24,25</sup>. On the other hand, considerable interest in the polyphenols present in plants and propolis has arisen with the recognition of their antioxidant effects.

A chemical analysis of propolis extracts revealed presence of flavones and flavonols<sup>1</sup>, which act by exerting direct positive inotropism<sup>26</sup>, but the prevailing belief is that they act by scavenging free radicals generated by doxorubicin and/or by chelating iron<sup>1,25,27,28</sup>. They act by (i) fixing of the free radicals on the DNA, (ii) activation of the detoxification

Table 2—Effect of doxorubicin (20 mg/kg), doxorubicin (20 mg/kg) + vinblastin (2 mg/kg) administrations alone or in association with the propolis extract on heart and liver GSH levels ( $\mu\text{g}/\text{mg}$  protein).[Values are mean  $\pm$  SD]

	Control n = 8	Doxorubicin (20mg/kg) n = 8	Doxorubicin (20mg/kg) + Vinblastin (2mg/kg) n = 5	Propolis (100mg/kg) + Doxorubicin (20mg/kg) n = 8	Propolis (100mg/kg) + Doxorubicin (20mg/kg) and Vinblastin (2mg/kg) n = 5
GSH in heart cell mitochondria	0.78 $\pm$ 0.06	0.33 $\pm$ 0.02*	0.68 $\pm$ 0.06*	0.83 $\pm$ 0.06	0.92 $\pm$ 0.07
GSH in liver cell mitochondria	1.10 $\pm$ 0.09	1.02 $\pm$ 0.09	0.62 $\pm$ 0.04*	1.21 $\pm$ 0.09	1.16 $\pm$ 0.10


\* $P < 0.001$ 

Fig. 6—Effect of doxorubicin (20 mg/kg), doxorubicin (20 mg/kg)+vinblastin (2 mg/kg) administrations alone or in association with the propolis extract on the liver mitochondrial SOD activity levels [i = doxorubicin (20 mg/kg), ii = doxorubicin (20 mg/kg)+vinblastin (2 mg/kg), iii = propolis+doxorubicin (20 mg/kg), iv = propolis+doxorubicin (20 mg/kg)+vinblastin (2 mg/kg)].




Fig. 7—Effect of doxorubicin (20 mg/kg), doxorubicin (20 mg/kg)+vinblastin (2 mg/kg) administrations alone or in association with the propolis extract on the heart and liver mitochondrial catalase activity levels [i = doxorubicin (20 mg/kg), ii = doxorubicin (20 mg/kg)+vinblastin (2 mg/kg), iii = propolis+doxorubicin (20 mg/kg), iv = propolis+doxorubicin (20 mg/kg)+vinblastin (2 mg/kg)].

system, (iii) protection of the lipid peroxidation, (iv) protection of cellular membranes<sup>14,15</sup>.

The antineoplastic drugs induced some transformations in the liver by the system cytochrome

P450. These transformations succeed to a highly toxic reactive metabolites formation for the cell<sup>29</sup> that interact with mitochondria leading to ROS production<sup>30,31</sup>. Most of chemotherapeutic agents like doxorubicin and cyclophosphamid are metabolized in the liver by the cytochrome P450<sup>29</sup> and the formed reactive intermediates are responsible for apoptotic process and cell toxicity<sup>32</sup>. The toxicity is function of the quantity of free radicals formed, the cellular reserve in GSH and the available glutathion-s-transférasés to assure the detoxification.

Doxorubicin and vinblastin altered mitochondrial functions as observed by a decrease in respiratory control value and the swelling activation, an SO overproduction and a lipid peroxidation either by heart and liver mitochondria. The data also showed mitochondrial functions restoration and polyphenols protective effect against the stress induced by chemotherapeutic agent. Polyphenols of propolis extract clearly reduced the drug toxic effect. Stress observed with the anticancer drug (decrease in lipid peroxidation, increase in GSH concentration and restoration of mitochondrial functions), was corrected entirely.

Treatment of Wistar rats with a single dose of 20 mg/kg/iv doxorubicin and 2 mg/kg vinblastin showed some effects in the heart and liver mitochondria functions especially the drugs decreases the RCR and increases SO and MDA productions. The doxorubicin has a moderate effect on the hepatic functions. In the group of animal's receiving 100 mg/kg propolis extract during 5 days before the treatment with chemotherapeutics drugs: restoration of the RCR, decrease of SO production and inhibition of the lipid peroxidation were observed. The same effects were observed *in vitro* on heart and liver mitochondria setting in presence of increasing concentrations of propolis extract. The results suggest prevention mechanism of propolis extract against doxorubicin cardiotoxicity and vinblastin hepatotoxicity. Doxorubicin and vinblastin are metabolized by the

liver to free radicals intermediates, which act on heart and liver mitochondria by increasing SO production, activation of membrane lipid peroxidation and therefore the fragilisation of the cell. Besides, the propolis inhibits the production of anion oxide by mitochondria. This effect is observed *in vivo* in the group of rats receiving the propolis orally before the administration of the chemotherapeutics agents whereas the rate of anion was increased significantly among the animals treated by only the drugs. The present results confirm the anti-oxidizer role of the flavonoids<sup>8,14</sup> which have the capacity to capture and deactivate the free radicals<sup>14,33,34,35</sup>.

A semiquinone radical is formed during liver doxorubicin metabolism<sup>36</sup>, suggesting that the mechanism of action of the propolis is similar in part to trans-resveratrol. The propolis polyphenols and trans-resveratrol, inhibit the  $O_2^{\cdot-}$  production in the cardiac and nervous cells, respectively<sup>37,38</sup>. This effect is due to the partial inhibition of the complex III (decylubiquinon cycle)<sup>39</sup>. The interaction of propolis polyphenols with the Q cycle inhibits the formation of radical ubisemiquinone; they can capture the first electron yielded by the cytochrome to the ubiquinol or can add an electron to the radical ubisemiquinone after departure of another of the ubiquinol in the Fe-S centre while inhibiting the formation of the same radical. The reduction of the radical  $O_2^{\cdot-}$  in the heart mitochondria can also be explained by the strong scavenger effect of the flavonoids as observed in the present study. Corroborating their antiradical properties, the flavonoids have direct actions against the radical  $OH\cdot$  hydroxyles and  $O_2^{\cdot-}$  superoxides that are species implied in the starting of the lipid peroxidation<sup>40</sup>. The inhibition of antioxidant system (GSH levels, catalase and SOD enzymes activities) by the anticancer drugs and their restoration by propolis extract reinforce this hypothesis. Alternatively, it is also possible as suggested by Koller<sup>41</sup> that the restoration of the mitochondrial membrane potential may be a way in the prevention of doxorubicin cardiotoxicity. A close relationship between the mitochondrial membrane potential and ROS production has been demonstrated<sup>4</sup>.

The present results indicate that propolis extract is able to restore the fall of membrane potential induced by doxorubicin and are able to protect the cell against the apoptosis of the drug.

## Conclusion

The results confirmed the role of propolis extract in protecting heart and liver tissues from doxorubicin and vinblastin toxicity by protecting the mitochondria

that should be given to minimise the damage membranes and enzymes. It is evident from the present work that the Algerian propolis is a very interesting antioxidant product that may be able to solve the problem of the anticancer drugs toxicity.

## References

- 1 Lahouel M, Boulkour S, Segueni N & Fillastre J P, Effet protecteur des flavonoides contre la toxicité de la vinblastine, du cyclophosphamide et du paracétamol par inhibition de la peroxydation lipidique et augmentation du glutathion hépatique, *Path Biol*, 52 (2004) 314.
- 2 Lahouel M & Fillastre J P, Role of flavonoïds in the prevention of hematotoxicity due to chemotherapeutic agents. *Haema*, 7 (2004) 313.
- 3 Sastre J, Pallardo F V, Garcia de la Asuncion J & Vina J, Mitochondria, oxidative stress and aging. *Free radic Res*, 32 (2000) 189.
- 4 Minotti G, Menna P, Salvatorelli E, Cairo G & Gianni L, Anthracyclines: Molecular advances and pharmacologic developments in antitumor activity and cardiotoxicity. *Pharmacol Rev*, 56 (2004) 185.
- 5 Andreeva L, Tanveer A & Crompton M, Evidence for the involvement of a membrane-associated cyclosporinA-binding protein in the  $Ca^{2+}$ - activated inner membrane pore of heart mitochondria, *Eur J Biochem*, 230 (1995) 1125.
- 6 Olson M & Kornbluth S, Mitochondria in Apoptosis and Human Disease, *Curr Mol Med*, 1 (2001) 91.
- 7 Pollack M & Leeuwenburgh C, Apoptosis and Aging: Role of the Mitochondria, *Gerontology*, 56 (2001) 475.
- 8 Harborne J B & Williams C A, Advances in flavonoid research since 1992, *Phytochem*, 55 (2000) 481.
- 9 Buratti S, Benedetti S & Cosio M S, Evaluation of the antioxidant power of honey, propolis and royal jelly by amperometric flow injection analysis, *Talanta*, 10 (2006) 1016.
- 10 Park Y K, Ikegaki M & Matias of Alencar S, Classificação das propolis Brasileiro a partir de suas características fisico-químicas e propriedades biológicas, *Mensagem Doce*, 58 (2000) 158.
- 11 Velikova M, Bankova V, Sorkun K, Houcine S, Tsetkova I & Kujemgiev A, Propolis from the mediterranean region : Chemical composition and antimicrobial activity, *Z Naturforsch*, 55 (2000) 790.
- 12 Serra Bonvehi J, Soliva Torrento´ M & Centelles Lorente E, Evaluation of Polyphenolic and Flavonoid Compounds in Honeybee-Collected Pollen Produced in Spain, *J Agric Food Chem*, 49 (2001) 1848.
- 13 Daneyemez M, Kurt E, Cosar A, Yuce E & Ide T, Methylprednisolone and vitamin E therapy in perinatal hypoxic-ischemic brain damage in rats, *Neuroscience*, 92 (1999) 693.
- 14 Siess M H, Le Bon A M, Canivenc-lavier M C & Suschetet M, Mechanisms involved in the chemoprevention of flavonoids, *Biofactors*, 12 (2000) 193.
- 15 Horakova K, Sovcikova A, Seemannova Z, Syrova D, Busanyova K, Drobna Z & Ferencik M, Detection of drug-induced, superoxide-mediated cell damage and its prevention by antioxidants, *Free Radic Biol Med*, 15 (2001) 650.
- 16 Morin D, Barthelemy S, Zini R, Labidalle S & Tillement J P,

- Curcumin induces the mitochondrial permeability transition pore mediated by membrane protein thiol oxidation, *Febs Letters*, 495 (2001a) 131.
- 17 Brooks P S, Digeress S B, Parks D A & Darley-Usmar V, Mitochondrial function in response to cardiac ischemia-reperfusion after oral treatment with quercetin, *Free Radic Biol Med*, 32 (2002) 1220.
- 18 Shen J G & Zhou D Y, Efficiency of Ginkgo biloba extract (EGB 761) in antioxidant protection against myocardial ischemia and reperfusion injury, *Biochem Biol*, 35 (1995) 125.
- 19 Zini R, Morin C, Bertelli A, Bertelli A A & Tillement J P, Effects of resveratrol on the rat respiratory chain, *Drugs Exp Clin Res*, 25 (1999) 87.
- 20 Bankova V & Marcucci M C, Standardization of propolis: present status and perspectives, *Bee World*, 81 (2000) 182.
- 21 Rustin P, Chrétien D, Bourgeron T, Gérard B, Rotig A, Saudubray J M & Munnich A, Biochemical and molecular investigations in respiratory chain deficiencies, *Clin Chim Acta*, 228 (1994) 35.
- 22 Clairborne A, Catalase activity. in: *CRC handbook of methods for oxygen radical research*, edited by RA Greenwald (CRC Press, Boca Raton) 1985, 288.
- 23 Beauchamp C & Fridovich I, Assay of superoxide dismutase, *Anal Biochem*, 44 (1971) 276.
- 24 Van Acker S A, Boven E, Van den Berg D J, Tromp M, Griffioen D H, van Bennekom W P, Van der Vijgh W J & Bast A, Structural aspect of antioxidant activity of flavonoids, *Free Rad Biol Med*, 20 (1996) 331.
- 25 Korshunov S S, Skulachev V P & Starkov A A, High protonic potential actuates a mechanism of production of reactive oxygen species in mitochondria, *FEBS Lett*, 416 (1997) 15.
- 26 Van Acker F A, Hulshof J W, Haenen G R, Menge W M, Van der Vijgh W J & Bast A, New synthetic flavonoids as potent protectors against doxorubicin-induced cardiotoxicity, *Free Radic Biol Med*, 31 (2001) 31.
- 27 Husken B C, De Jong J, Beekman B, Onderwater R C, Van der Vijgh W J & Bast A, Modulation of the *in vitro* cardiotoxicity of doxorubicin by flavonoids, *Cancer Chemother Pharmacol*, 37 (1997) 55.
- 28 Van Acker S A, Boven E, Kuiper K, Van den Berg D J, Grimbergen J A, Kramer K, Bast A & Van der Vijgh W J, Monohydroxyethylrutoside, a dose-dependent cardioprotective agent, does not affect the antitumor activity of doxorubicin, *Clin Cancer Res*, 3 (1997) 1747.
- 29 Grattagliano I, Portincasa P, Palmieri V O & Palasciano G, Overview on the mechanisms of drug-induced liver cell death, *Annals Hepatology*, 1 (2002) 162.
- 30 Lemasters J J, Nieminen A L, Qian T, Trost L C, Elmore S P, Nishimura Y & al., The mitochondrial permeability transition in cell death: a common mechanism in necrosis, apoptosis and autophagy, *Biochem Biophys Acta*, 1366 (1998) 177.
- 31 Morin D, Hauet T, Spedding M & Tillement J P, Mitochondria as a target for antiischemic drugs, *Adv Drug Deliv Rev*, 1 (2001b) 1254.
- 32 Green D R & Kroemer G, The pathophysiology of mitochondrial cell death, *Science*, 305 (2004) 626.
- 33 Cotelle N, Role of flavonoids in oxidative stress, *Curr Top Med Chem*, 6 (2001) 569.
- 34 Ichikawa H, Satoh K, Tobe T, Yasuda I, Ushio F, Matsumoto K & al., Free radical scavenging activity of propolis, *Redox Rep*, 7 (2002) 347.
- 35 Silva M M, Santos M R, Carço G, Rocha R, Justino G & Mira L, Structure –antioxidant activity relationships of flavonoids: a re-examination, *Free Radic Res*, 36 (2002) 1219.
- 36 De Matteis F, The role of cytochrome P<sub>450</sub> in drug metabolism and toxicity, (CRC press, New York) (1994), 81.
- 37 Kamei H, Koide T, Kojimam T & Hasegawa M, Flavonoid-mediated tumor growth suppression demonstrated by *in vivo* study, *Cancer Biother Radiophar*, 3 (1996) 193.
- 38 Zini R, Simon N, Morin C, D'Athis P & Tillement J P, Inhibition of rat cerebral mitochondrial respiration by cyclosporins A, D, and G and restoration with trimetazidine, *Paris CR, Acad Sci*, 319 (1996) 1087.
- 39 Degli-Esposti M, Inhibitors of NADH-ubiquinone reductase: an overview, *Biochim Biophys Acta*, 1364 (1998) 222.
- 40 Moreno A J, Santos D J & Palmeira C M, Ischemic heart disease: the role of mitochondria-carvedilol prevents lipid peroxidation of mitochondrial membranes, *Rev Port Cardiol*, 17 (1998) 1163.
- 41 Koller A & Haid C, Is restoring regenerative potential a way to prevent anthracycline cardiotoxicity? A hypothesis, *Med Sci Monit*, 11 (2005) 21.