

Acid and salt tolerance behavior of *Rhizobium* isolates and their effect on microbial diversity in the rhizosphere of redgram (*Cajanus cajan* L.)

Debadatta Sethi*, Santanu Mohanty & Sushanta Kumar Pattanayak

Department of Soil Science and Agricultural Chemistry, Odisha University of Agriculture and Technology,
Bhubaneswar- 751 003, Odisha, India

Received 10 January 2019; revised 04 February 2019

Experiments were conducted in two phases, first under *in vitro* condition to study the stress tolerant ability and then in pot experiment to study the effect of *Rhizobium* isolates on rhizospheric microbial activity. The strain CHRS-7 could tolerate the pH 4.0, whereas RAN-1 and RAB-1 could not. The growth of all the strains was luxuriant in 1% NaCl solution and decreased with increase in the concentration of NaCl. All the strains could produce the phytohormone indole acetic acid (IAA) by metabolizing different carbon sources. The highest amount of IAA was produced by RAB-1 (81 µg/mL), CHRS-7 (78 µg/mL) and RAN-1 (72 µg/mL) by metabolizing mannitol, glucose, and sucrose, respectively. The higher bacteria and rhizobial population was enumerated in the treatment with inoculation of *Rhizobium* strains and added with 50% of soil test dose of nitrogen whereas higher fungi population was enumerated with the treatment receiving 150% of a soil test dose of nitrogen. The soil enzymes activity, microbial biomass carbon and nitrogen were also higher with a lower dose of external sources of N (50% of a soil test dose) and decreased with increase in nitrogen dose.

Keywords: *Cajanus cajan* L., Microbial biomass carbon (MBC), Microbial biomass nitrogen (MBN), Plant growth promoting rhizobacteria (PGPR), Root nodule, Stress tolerance

The plant growth promoting rhizobacteria including symbiotic nitrogen (N₂) fixers are important for the sustainability of agriculture¹⁻⁴. Approximately 2.5×10^{11} kg NH₃ yr⁻¹ is fixed from the atmosphere through biological nitrogen fixation (BNF) by legumes and cyanobacteria⁵. The adverse environmental factors, like soil acidity⁶, and soil salinity⁷ influence symbiotic nitrogen fixation. Superior growth of *Rhizobium* has been reported at neutral pH (7.0). The soil acidity limits *Rhizobium* survival, persistence, and nodulation⁸ hampering symbiotic nitrogen fixation in both tropical and temperate soils. The Bradyrhizobia have generally been considered to be tolerance to acidic pH^{9,10}. Different mechanisms are involved for adapting to acidic environments, including the use of proton pump systems^{11,12} or glutamic acid decarboxylase¹³ for increasing the internal pH.

Acidic soil does not allow the rhizobial cells to survive in adequate numbers in free-living state. The rhizobial strains of a given species vary widely in their pH tolerance¹⁴. The optimum pH range for growth of rhizobia was between 5.5-7.5¹⁵. The acid tolerant strains were able to grow at pH 4.5 to 5.5¹⁶.

The ability to grow at an acidic pH would provide these isolates with a competitive advantage over another rhizosphere organisms¹⁰.

The salinity of the soil develops due to a nutrient imbalance in the soil, which is considered as a constraint influencing the N₂ fixing symbiosis and the survival of both partners^{7,17}. The symbiotic rhizobia are more sensitive to salt stress than free-living rhizobia¹⁸. Some species of rhizobia adapt to saline conditions through the intracellular accumulation of low-molecular-weight organic solutes called osmolytes, such as glutamate, trehalose, glycine betaine and polyamines, or accumulation of potassium ion (K⁺)¹⁹. The growth of isolate *Sinorhizobium meliloti* was not completely inhibited by 5% NaCl^{20,21}. There have been several studies and the isolates have shown the disparity in tolerance to different concentration of salt²².

The plant hormone production in nodules by microsymbiont and its transport to host by nodular symbiont was first reported by Hunter²³ in soybean. The PGPR effect of rhizobia creates a better environment not only for their own activity but also for other microbes in both legume and nonlegume rhizosphere. The IAA is one of the most physiologically active auxins. It is a common

*Correspondence:
E-mail: debadattaouat@gmail.com

product of L-tryptophan metabolism by several microorganisms^{24,25}. The amount of IAA produced varied from strain to strain²⁶. The rhizobia isolated from the root nodules of *Cajanus cajan*²⁷, *Dalbergia lanceolaria*²⁸, *Roystonea regia*²⁹ produced high amounts of IAA during growth in basal medium supplemented with L-tryptophan. The maximum amount of IAA was produced when mannitol was used as carbon source in the isolate from *I. viscosa*³⁰, *Cajanus cajan*²⁷, and glucose in *Vigna mungo*³¹. Other rhizobacteria like *Pseudomonas fluorescens* produced IAA by utilizing tryptophan³².

Besides nodulation and nitrogen fixation, seed inoculation with rhizobia can stimulate the production of phytohormones, siderophores, and hydrogen cyanide as well as soil microbial diversity and structure, potentially enhancing PGPR activity^{33,34}. Inoculation with an *R. gallicum* strain induced growth of bacterial communities that had been often reported as PGPM (plant growth-promoting microorganisms)³³.

The release of bioinoculants in huge amount may either result in non-target effects, which in turn enhance plant growth or lead to an ecological risk³⁵. The rhizobia-legume interaction is assumed to act as a driving force for maintaining the nitrogen balance in the soil. This effect also depends on the rhizosphere and internal nitrogen turnover³⁶. It is therefore important to determine the effect of rhizobial inoculants on the resident microbial community. The *Bradyrhizobium* sp. inoculation has positively affected the population of actinomycetes and pseudomonas during the early stages of *Cajanus cajan* growth³⁷. The enzyme (dehydrogenases, urease, protease, and acid phosphomonoesterase) activities enhanced due to inoculation of *R. leguminosarum* bv. *Viciae* in the rhizosphere of *Vicia faba* L.³⁸

The objective of the present investigation was to isolate an acid and salinity tolerant strain of *Rhizobium* for red gram crop and to find out their effect on microbial diversity and enzyme activity in redgram rhizosphere.

Materials and Methods

Indole-3-acetic acid (IAA) production

The IAA was obtained from 6 days old cultures of *Rhizobium* spp. incubated at 30°C and 150 rpm in YEM broth with 0.1 g/L tryptophan. After centrifugation (6000 rpm/30 min), the liquid portion of an aliquot of the broth was used to determine the concentration of indole-3-acetic acid by the method described by Glickman and Dessaux³⁹.

Effect of pH and salt concentration on growth of isolate

The isolates were inoculated to nutrient broth medium maintained at pH 4.0, 5.0, 6.0, 7.0 and 8.0 to measure acid tolerance and for salinity tolerance, NaCl was added @ 1%, 2%, 3%, 4%, and 5%, respectively and incubated at 30°C for 96 h. The turbidity of the medium in each flask was measured at 660 nm using visible spectrophotometer at 3 h interval till 24 h, 6 h interval till 36 h and 12 h interval till 96 h of inoculation.

Pot experiment

Characterization of experimental soil

The soil was collected from the experimental plot of the horticultural research station of Odisha University of Agriculture and Technology, Bhubaneswar and processed for the physicochemical and biological properties estimation, by following different standard methods.

Microbial population was estimated by serial dilution and spread plate technique⁴⁰ by using growth media *viz*: nutrient agar, rosebengal chromaphenicol agar and yeast mannitol agar (Himedia) for total heterotrophic bacteria, fungi and *Rhizobium*, respectively. The dehydrogenase activity⁴¹, Urease activity⁴², chloroform fumigation extracted microbial biomass carbon⁴³ and microbial biomass nitrogen⁴⁴ were determined by standard methods.

Preparation of pots for sowing

Five kilograms of processed soil was filled in each pot. Red gram cv Asha seeds were surface sterilized by 4% sodium hypochlorite and ethyl alcohol, washed by sterilized distilled water and inoculated with a liquid culture of strain CHRS-7, RAN-1 and RAB-1 separately for 30 min. The inoculated seeds were sown at 2 cm depth of soil in the pot and after germination one plant was maintained in each pot. The pest and disease infestation was monitored and care was taken accordingly. After the 90th day of sowing, the plants were uprooted and the rhizospheric soils were collected for estimating microbial parameters.

Treatment combinations

The three rhizobium species were inoculated to the seeds which received four different doses of soil test based nitrogen (0, 50, 100 and 150 %) *viz*: T₁-CHRS7 + N₀, T₂-CHRS7 + N₅₀, T₃-CHRS7 + N₁₀₀, T₄-CHRS7 + N₁₅₀, T₅-RAN-1 + N₀, T₆-RAN-1 + N₅₀, T₇-RAN-1 + N₁₀₀, T₈-RAN-1 + N₁₅₀, T₉-RAB-1 + N₀, T₁₀-RAB-1 + N₅₀, T₁₁-RAB-1 + N₁₀₀ and T₁₂-RAB-1 + N₁₅₀. The

phosphorus and potassium were applied as per soil test dose to all the treatments. Each treatment was replicated thrice and imposed for factorial CRD (completely randomized design). The test crop was redgram (*Cajanus cajan*) cv: ASHA (ICPL 87119).

Statistical analysis

The data was analyzed statistically as per the procedure prescribed for complete randomized design for the *in vitro* experiment and factorial CRD for pot experiment by using the SPSS software.

Result and Discussion

Total 56 numbers of isolates were selected for biochemical characterization (data not given), out of which only twelve isolates were screened out for the molecular characterization and after molecular characterization three isolates (CHRS-7, RAN-1, and RAB-1) were confirmed as *Rhizobium* spp. The 16S rRNA sequences of the three isolates were submitted to NCBI and assigned as accession number MH636329, MH541051 and MH636773 for CHRS-7, RAN-1 and RAB-1, respectively.

The production of IAA by three isolates:-

All the three isolates were grown in mineral medium containing eight types of carbon sources (mannitol, glucose, sucrose, galactose, starch, fructose, and maltose) at 1% concentration and IAA production was estimated (Fig. 1). The strain RAB-1 metabolized mannitol to produce highest (81 µg/mL) amount of IAA followed by glucose (62 µg/mL), fructose (45 µg/mL), starch (42 µg/mL), lactose (39 µg/mL), maltose (34 µg/mL), sucrose (30 µg/mL) and galactose (21 µg/mL). The lowest was estimated in control (17 µg/mL). The strain CHRS-7 was able to metabolize glucose to produce maximum (78 µg/mL) amount of IAA followed by fructose (60 µg/mL), mannitol (55 µg/mL), sucrose (42 µg/mL), starch (39 µg/mL), lactose (35 µg/mL), galactose (23 µg/mL) and maltose (22 µg/mL) and the least (16 µg/mL) was recorded in control. The strain




Fig. 1 — The production of Indole acitic acid (IAA) by the three isolates

RAN-1 metabolized sucrose to produce maximum (72 µg/mL) amount of IAA followed by fructose (51 µg/mL), starch and glucose (45 µg/mL), maltose (37 µg/mL), mannitol (36 µg/mL), lactose (28 µg/mL) and galactose (27 µg/mL). The least (21 µg/mL) was recorded in control. The utilization of carbon sources by the rhizobia for production of IAA corroborate the findings of Kumar & Ram⁴⁶, Sarkar & Laha⁴⁷, Ghosh *et al.*⁴⁸, Kucuk and Cevheri⁴⁹. The production of IAA by utilization of tryptophan was reported in *Pseudomonas fluorescens*³².

Acid and salt tolerance capacity of isolates

The acid tolerance capacity of the three isolates was determined by measuring their growth in YEM broth expressed as optical density (OD) at different pH *viz* 4.0, 5.0, 6.0, 7.0 and 8.0 (Fig. 2A-E). In the growth media of pH 4.0, the OD of strain CHRS-7 was increasing up to 48 h and then attained the plateau stage (OD 1.5) until 96 h. Other two isolates (RAN-1 and RAB-1) were unable to grow in the media and the OD was below 0.5 till 96 h. A similar trend of OD was observed in the media of pH 5.0. At pH 6.0 the OD of RAN-1 and RAB-1 were recorded less than that of CHRS-7 up to 30 h. After 36 h of growth, all the OD became stagnant till 96 h. The




Fig. 2 — Growth of isolates in YEM broth of (A) pH-4.0; (B) pH-5.0; (C) pH-6.0; (D) pH-7.0; and (E) pH-8.0

maximum OD of all the three isolates was below 2.0. At pH 7.0, the OD for CHRS-7 and RAN-1 were above 2.0, whereas, OD for RAB-1 was below 2.0 till 96 h. At pH 8.0 all the three isolates OD recorded ≥ 2.0 till 96 h. Similar results on acid tolerance of rhizobia were reported by Shamseldin & Werner⁵⁰, Belal *et al.*⁵¹, and Guerrero-Castro *et al.*⁵².

In 1% NaCl medium, all the strains grew luxuriantly up to 51 h; thereafter the growth was stagnant (Fig. 3A). In 2% solution, the growth of strains was higher up to 51 h, and stagnant thereafter. The growth of RAN-1 was higher among the three isolates till 99 h (Fig. 3B). Similar trend was recorded with 3 % NaCl (Fig. 3C). In 4 % NaCl, the stationary phase of RAB-1 and CHRS-7 was observed after 21 h, whereas stationary phase of RAN-1 was observed after 51 h (Fig. 3D). The OD of CHRS-7 and RAB-1 was below 0.5 till 99 h, whereas, the OD of RAN-1 was 1.83 after 51 h and maintained till 99 h. In 5 % NaCl the stationary phase came after 21 h and the OD of all the three strains were below 0.5 (Fig. 3E). The findings were in agreement with Omotoyinbo & Omotoyinbo⁵³ in *Escherichia coli*, Kucuk *et al.*⁴⁹ in *Staphylococcus aureus*, Rai *et al.*⁵⁴, and Patil *et al.*⁵⁵ in *Rhizobium* sp.

Pot experiment

Initial physico-chemical and biological properties of experimental soil.

The experimental soil was sandy loam in texture with bulk density 1.6 Mg/m³, particle density 2.45 Mg/m³

and porosity of 34.7%. The soil was acidic in reaction (pH-4.92) with organic carbon 05.4 g kg⁻¹ soil and low in KMnO₄ mineralizable N (199 kg ha⁻¹). The heterotrophic bacteria, fungi and rhizobia population were 2.1 × 10⁶ cfu g⁻¹, 1.7 × 10⁵ cfu g⁻¹ and 1.9 × 10⁵ cfu g⁻¹ soil, respectively. The dehydrogenase and urease activities, microbial biomass carbon (MBC) and microbial biomass nitrogen (MBN) were 18 µg TPF g⁻¹dw, 2.3 µg of NH₄⁺ g⁻¹ soil h⁻¹, 110 µg C kg⁻¹ soil and 18.8 µg N g⁻¹ dry soil, respectively.

Effect of rhizobial inoculation on microbial diversity

The data related to total heterotrophic bacteria, fungi and rhizobial population have been present in (Table 1).

The total bacterial population in initial soil was 2.1 × 10⁶ cfu g⁻¹ and after 90 days after sowing (DAS) it ranged from 3.0 × 10⁶ cfu g⁻¹ to 6.8 × 10⁶cfu g⁻¹. The strain CHRS-7 inoculated pot had more bacterial population than those inoculated with strains RAN-1 and RAB-1. The maximum bacterial population (6.8 × 10⁶ cfu g⁻¹) was estimated in the pot where strain CHRS-7 was inoculated along with 50% of soil test dose (STD) N. Irrespective of the strains the bacterial population increased upto 50% of STD N and thereafter decreased with increase in nitrogen concentration. The initial fungi population was 1.7 × 10⁵ cfu g⁻¹. The fungal population at 90 DAS ranged from 4.7 × 10⁵ cfu g⁻¹ to 9.7 × 10⁵ cfu g⁻¹. The results revealed that the increase in the concentration




Fig. 3 — A, B, C, D and E indicate the growth curves of isolates in 1%, 2%, 3%, 4%, and 5% NaCl solution, respectively

Table 1 — Effect of inoculation of *Rhizobium* spp. on microbial diversity

Effect of rhizobia inoculation on bacterial population ($\times 10^6$ cfu g^{-1} soil)					
	N ₀	N ₅₀	N ₁₀₀	N ₁₅₀	Mean-S
CHRS-7	3.9	6.8	6.6	4.7	5.5
RAN-1	3.5	4.8	4.8	3.5	4.2
RAB-1	3.0	4.7	4.6	3.7	4.0
Mean-N	3.5	5.4	5.4	4.0	
LSD ($P=0.05$) S (0.2) N (0.3) S×N (0.5)					

Effect of rhizobia inoculation on fungal population ($\times 10^5$ cfu g^{-1} soil)

	N ₀	N ₅₀	N ₁₀₀	N ₁₅₀	Mean-S
CHRS-7	5.6	8.1	8.3	9.7	7.9
RAN-1	5.2	6.8	7.1	8.7	7.0
RAB-1	4.7	6.2	6.1	8.1	6.3
Mean-N	5.2	7.0	7.2	8.8	
LSD ($P=0.05$) S (0.4) N (0.5) S×N (0.9)					

Effect of rhizobia inoculation on rhizobial population ($\times 10^5$ cfu g^{-1} soil)

	N ₀	N ₅₀	N ₁₀₀	N ₁₅₀	Mean-S
CHRS-7	5.0	8.1	7.8	6.0	6.7
RAN-1	4.6	6.1	6.0	4.8	5.4
RAB-1	4.1	5.9	5.9	5.0	5.2
Mean-N	4.6	6.7	6.6	5.3	
LSD ($P=0.05$) S (0.4) N (0.6) S×N (0.7)					

of nitrogenous fertilizer increased the fungal population. The initial rhizobial population was 1.9×10^5 cfu g^{-1} and at 90 DAS it increased in all the treatments ranging from 3.0×10^5 cfu g^{-1} to 6.8×10^5 cfu g^{-1} soil. The highest rhizobial population was enumerated in the treatment where the strain CHRS-7 was inoculated with 50% of a soil test dose of nitrogen and thereafter it decreased. The rhizobial communities in the rhizosphere were inhibited by nitrogen fertilization, which may be related to previous reports on the negative effect of nitrogen fertilization on nodulation³². The better growth of plants in inoculation of rhizobia could improve the quality and quantity of root exudates and the incorporation of root biomass to the soil, which subsequently benefited the microbial growth in the rhizosphere⁵⁶⁻⁵⁸.

Effect of rhizobial inoculation on enzyme activity

The dehydrogenase and urease activities of initial soil were $18 \mu\text{g TPF } g^{-1}$ and $2.3 \mu\text{g of } NH_4^+ g^{-1} \text{ soil } h^{-1}$, respectively. The dehydrogenase activity at 90 DAS ranged from $38 \mu\text{g TPF } g^{-1} \text{ dw}$ to $85 \mu\text{g TPF } g^{-1} \text{ dw}$ and urease activity ranged from $6 \mu\text{g of } NH_4^+ g^{-1} \text{ soil } h^{-1}$ to $14 \mu\text{g of } NH_4^+ g^{-1} \text{ soil } h^{-1}$ (Table 2). The highest

Table 2 — Effect of inoculation of *Rhizobium* spp. on enzyme activity

Effect of rhizobia inoculation on Dehydrogenase ($\mu\text{g TPF } g^{-1} \text{ dw}$)					
	N ₀	N ₅₀	N ₁₀₀	N ₁₅₀	Mean-S
CHRS-7	59	85	75	56	70
RAN-1	39	58	53	37	47
RAB-1	38	55	48	34	44
Mean-N	45	66	59	42	
LSD ($P=0.05$) S (3.7) N (4.3) S×N (NS)					

Effect of rhizobia inoculation on urease ($\mu\text{g of } NH_4^+ g^{-1} \text{ soil } h^{-1}$) activity

	N ₀	N ₅₀	N ₁₀₀	N ₁₅₀	Mean-S
CHRS-7	10	14	12	10	11
RAN-1	6	10	8	7	8
RAB-1	5	8	6	5	6
Mean-N	7	11	9	7	
LSD ($P=0.05$) S (1.4) N (1.6) S×N (NS)					

dehydrogenase activity and urease activity were estimated in the treatment where strain CHRS-7 was inoculated and the crop received 50% nitrogen of STD. followed by N₁₀₀, N₀, and N₁₅₀. In higher nitrogen concentration, the enzyme activity reduced because of the reduced population of bacteria. The findings are agreement with Gopalakrishnan *et al.*⁶², Siczek and Lipiec³⁸ and Fall *et al.*⁶⁰. They reported that the rhizobial inoculation increased total microbial activities and acid phosphatase activity.

Effect of rhizobia inoculation on microbial biomass 'C' and 'N'

The MBC reflects the size of the soil microbial community and is believed to be an indicator of microbiological properties and soil fertility⁵⁸. The microbial biomass carbon and nitrogen of initial soil was $98 \mu\text{g C } kg^{-1}$ soil and $18.8 \mu\text{g N } g^{-1}$, respectively. The highest microbial biomass carbon ($268 \mu\text{g C } kg^{-1}$ soil) and nitrogen ($59.8 \mu\text{g N } g^{-1}$ dry soil) were estimated in the treatment where *Rhizobium* strain CHRS-7 was inoculated with the application of 50% soil test dose N (Table 3). The lowest microbial biomass carbon ($108 \mu\text{g C } kg^{-1}$ soil) and nitrogen ($238 \mu\text{g N } g^{-1}$ dry soil) were estimated in the treatment where strain RAB-1 was inoculated with application 150% soil test dose N. With the application of higher dose of nitrogen the microbial biomass carbon and nitrogen reduced due to inhibitory influence of N on bacterial community in the rhizosphere^{60,61}.

Relationship of a bacterial population with urease, dehydrogenase, MBC, and MBN

The relation of the bacterial population with MBC, MBN and enzyme activity were presented in (Fig. 4).

It indicated that the rhizospheric bacterial population was positively correlated with urease activity, dehydrogenase activity; microbial biomass carbon and microbial biomass nitrogen with a coefficient of determination is 0.751, 0.858, 0.864 and 0.846, respectively. The coefficient of determination (R^2) indicated that the increase of bacterial population

enhanced the enzyme activity, microbial biomass carbon, and nitrogen.

Conclusion

This study concluded that the microsymbiont (CHRS-7) could tolerate the pH up to 4.0, whereas others couldn't. All the microsymbionts were able to grow better in 1% NaCl concentration and were inhibited with an increase in concentration of NaCl. Both the total heterotrophic bacteria and rhizobia population were higher in suboptimal dose of N (50% of soil test dose), whereas fungal population was higher in an optimal dose of N (100% of soil test dose) as well as in super optimal dose of N (150% of soil test dose). The enzyme activity and microbial biomass carbon and nitrogen were higher in suboptimal dose and decreased with an increase in dose of nitrogen. The bacterial population in rhizosphere had a positive effect on enzyme activity, MBC, and MBN.

Table 3 — Effect of rhizobia inoculation on microbial biomass carbon and nitrogen

Effect of rhizobia inoculation on microbial biomass carbon (MBC) ($\mu\text{g C kg}^{-1}$ soil)

	N ₀	N ₅₀	N ₁₀₀	N ₁₅₀	Mean-S
CHRS-7	180	268	229	186	216
RAN-1	125	174	165	119	146
RAB-1	115	162	153	108	134
Mean-N	140	201	182	138	
LSD ($P=0.05$)	S (5.4)	N (6.2)	S×N (10.8)		

Effect of rhizobia inoculation on microbial biomass nitrogen (MBN) ($\mu\text{g N g}^{-1}$ dry soil)

	N ₀	N ₅₀	N ₁₀₀	N ₁₅₀	Mean-S
CHRS-7	40	59	51	42	48
RAN-1	27	40	36	25	32
RAB-1	26	38	33	23	30
Mean-N	31	45	40	30	
LSD ($P=0.05$)	S (5.6)	N (6.5)	S×N (NS)		

Acknowledgment

The authors acknowledge the financial support of All India Network Project on Soil Biodiversity-Biofertilizer (AINP on SBB) of Indian Council of Agriculture Research (ICAR).




Fig. 4 — (A) Relationship between bacterial population and urease activity (B) Dehydrogenase activity; (C) Microbial biomass carbon (MBC); and (D) and Microbial biomass nitrogen (MBN)

References

- 1 Barea JM, Future challenges and perspectives for applying microbial biotechnology in sustainable agriculture based on a better understanding of plant-microbiome interactions. *J Soil Sci Plant Nutr*, 15 (2015) 261.
- 2 Lagos L, Maruyama F, Nannipieri P, Mora ML, Ogram A & Jorquera MA, Current overview on the study of bacteria in the rhizosphere by modern molecular techniques: a mini-review. *J Soil Sci Plant Nutr*, 15(2015) 504.
- 3 Larsen J, Jaramillo-López P, Nájera-Rincon M & González-Esquivel CE, Biotic interactions in the rhizosphere in relation to plant and soil nutrient dynamics. *J Soil Sci Plant Nutr*, 15 (2015) 449.
- 4 Oliveira DP, de Figueiredo MA, Soares BL, Teixeira OHS, Martins FAD, Rufini M, Chain CP, Reis RP, de Moraes AR, de Souza FMM & de Andrade MJB, Acid tolerant *Rhizobium* strains contribute to increasing the yield and profitability of common bean in tropical soils. *J Soil Sci Plant Nutr*, 17 (2017) 922.
- 5 Pattanayak SK, Biological Nitrogen Fixation – Status, Potential and Prospects in Supplementing Nitrogen Needs of the Crops. *Indian J Fert*, 12(2016) 94.
- 6 Vinuesa P, Neumann-Silkow F, Pacios-Bras C, Spaink HP, Martínez-Romero E & Werner D, Genetic analysis of a pH-regulated operon from *Rhizobium tropici* CIAT899 involved in acid tolerance and nodulation competitiveness. *Mol Plant Microbe Interact*, 16 (2003) 159.
- 7 Mohammadi K, Sohrabi Y, Heidari G, Khaledi S & Majidi M, Effective factors on biological nitrogen fixation. *Afr J Agric Res*, 7 (2012) 1782.
- 8 Brockwell J, Bottomley PJ & Thies JE, Manipulation of rhizobia microflora for improving legume productivity and soil fertility: A critical assessment. *Plant Soil*, 174 (1995) 143.
- 9 Appunu C & Dhar B, Symbiotic effectiveness of acid-tolerant *Bradyrhizobium* strains with soybean in low pH soil. *Afr J Biotechnol*, 5 (2006) 842.
- 10 Appunu C, Reddy LML, Reddy CVC, Sen D & Dhar B, Symbiotic diversity among acid-tolerant bradyrhizobial isolates with cowpea. *J Agric Sci*, 4 (2009) 126.
- 11 Papadimitriou K, Pratsinis H, Nebe-von-Caron G, Kletsas D & Tsakalidou E, Acid tolerance of *Streptococcus macedonicus* as assessed by flow cytometry and single-cell sorting. *Appl Environ Microbiol*, 73 (2007) 465.
- 12 Lei Z, Jian-ping G, Shi-qing W, Ze-yang Z, Chao Z & Yongxiong Y, Mechanism of acid tolerance in a *rhizobium* strain isolated from *Pueraria lobata* (Willd.) Ohwi. *Can J Microbiol*, 57 (2011) 514.
- 13 Gandhi M & Chikindas ML, Listeria: a food borne pathogen that knows how to survive. *Int J Food Microbiol*, 113 (2007) 1.
- 14 Appunu C, Sen D & Dhar B, Acid and aluminium tolerance of *Bradyrhizobium* isolates from traditional soybean growing areas of India. *Indian J Agric Sci*, 75 (2005) 727.
- 15 Gauri, Singh AK, Bhatt RP, Pant S, Bedi MK & Naglot A, Characterization of *Rhizobium* isolated from root nodules of *Trifolium alexandrinum*. *J Agric Technol*, 7 (2011) 1705.
- 16 Keneni A, Assefa F & Prabu PC, Characterization of acid and salt tolerant rhizobial strains isolated from faba bean fields of wollo, northern ethiopia. *J Agric Sci Tech*, 12 (2010) 365.
- 17 Niste M, Vidican R, Pop R & Rotar I, Stress factors affecting symbiosis activity and nitrogen fixation by *Rhizobium* cultured *in vitro*. *Pro Environ*, 6 (2013) 42.
- 18 Zahran HH, *Rhizobium*-legume symbiosis and nitrogen fixation under severe conditions and in an arid climate. *Microbiol Mol Biol Rev*, 63 (1999) 968.
- 19 Wei W, Jiang J & Yang SS, Mutagenesis and Complementation of rel A from *Sinorhizobium meliloti* 042BM as a Salt Tolerance Involvement Gene. *Annals Microbiol*, 54 (2004) 317.
- 20 Saraf M & Dhandhukia P, Response of *Sinorhizobium meliloti* to High Salt Concentration and Effect of Added Osmotica. *J Microb World*, 7 (2005) 250.
- 21 Kucuk CM, Kivanç M & Kinaci E, Characterization of *Rhizobium* sp. Isolated from Bean. *Turk J Biol*, 30 (2006) 127.
- 22 Fentahun M, Akhtar M, Muleta D and Lemessa F, Isolation and characterization of nitrogen deficit *Rhizobium* isolates and their effect on growth of haricot bean. *Afr J Agric Res*, 8 (2013) 5942.
- 23 Hunter WJ, Indole-3-Acetic acid production by bacteroids from soybean root nodules. *Physiol Plant*, 76 (1989) 31.
- 24 Ghosh S & Basu PS, Production and metabolism of Indole acetic acid in roots and root nodules of *Phaseolus mungo*. *Microbiol Res*, 161 (2006) 362.
- 25 Mandal SM, Mondal KC, Dey S & Pati BR, Optimization of cultural and nutritional conditions for indole-3-acetic acid (IAA) production by a *Rhizobium* sp. isolated from root nodules of *Vigna mungo* (L.) Hepper. *Res J Microbiol*, 2 (2007) 239.
- 26 Garg A & Sharma M, Evaluation of phosphate solubilizing activity and indole acetic acid production of rhizobia. *Int J Sci Res*, 2 (2013) 314.
- 27 Shende RC & Patil MB, Growth behaviour and Indole Acetic Acid (IAA) production by a *Rhizobium* sp. Isolated from *Cajanus cajan* plant. *Int J Pharm Biol Sci*, 2 (2011) 621.
- 28 Ghosh AC & Basu PS, Growth behavior and bio-reduction of indole acetic acid by a *Rhizobium* sp. isolated from root nodules of a leguminous tree *Dalbergia lanceolaria*. *Indian J Exp Biol*, 40 (2002) 796.
- 29 Basu PS & Ghosh AC, Production of indole acetic acid in culture by a *Rhizobium* species from the root nodules of a monocotyledonous Tree, *Roystonea regia*. *Acta Biotechnol*, 21 (2001) 65.
- 30 Kumari BS, Ram MR & Mallaiah KV, Studies on exopolysaccharide and indole acetic acid production by *Rhizobium* strains from Indigofera. *Afr J Microbiol Res*, 3 (2009) 10.
- 31 Satyanandam T, Babu K, Rosaiah G & Vijayalakshmi M, Screening of *Rhizobium* strains isolated from *Vigna mungo* native to rice fallows for the production of indole acetic acid. *Int J Pharm Biol Sci*, 3 (2013) 56.
- 32 Saxena J, Saini A, Kushwaha K & Ariño A, Synergistic effect of plant growth promoting bacterium *Pseudomonas fluorescens* and phosphate solubilizing fungus *Aspergillus awamori* for growth enhancement of chickpea. *Indian J Biochem Biophys*, 53(2016) 135.
- 33 Trabelsi D, Mengoni A, Ammar HB & Mhamdi R, Effect of on-field inoculation of *Phaseolus vulgaris* with rhizobia on soil bacterial communities. *FEMS Microbiol Ecol*, 77 (2011) 211.
- 34 Trabelsi D, Ammar HB, Mengoni A & Mhamdi R, Appraisal of the crop-rotation effect of rhizobial inoculation on potato cropping systems in relation to soil bacterial communities. *Soil Biol Biochem*, 54 (2012) 1.

- 35 Gupta R, Bru D, Bisaria VS, Philippot L & Sharma S, Responses of *Cajanus cajan* and rhizospheric N-cycling communities to bioinoculants. *Plant Soil*, 358 (2012) 143.
- 36 Babic KH, Schauss K, Hai B, Sikora S, Reazepovic S, Radl V & Schloter M, Influence of different *Sinorhizobium meliloti* inocula on abundance of genes involved in nitrogen transformations in the rhizosphere of alfalfa (*Medicago sativa* L.). *Environ Microbiol*, 10 (2008) 2922.
- 37 Sharma R, Paliwal JS, Chopra P, Dogra D, Pooniya V, Bisaria VS, Swarnalakshmi K & Sharma S, Survival, efficacy and risk assessment of bacterial inoculants in *Cajanus cajan*. *Agric Ecosyst Environ*, 240 (2017) 244.
- 38 Siczek A & Lipiec J, Impact of faba bean-seed rhizobial inoculation on microbial activity in the rhizosphere soil during growing season. *Int J Mol Sci*, 17 (2016) 1.
- 39 Glickman M & Dessaux Y, A critical examination of the specificity of the salkowski reagent for indolic compounds produced by phytopathogenic bacteria. *Appl Environ Microbiol*, 61(1995) 793.
- 40 Ben-David A & Davidson CE, Estimation method for serial dilution experiments. *J Microbiol Methods*, 117 (2014) 212.
- 41 Tabatabai MA, *Soil enzymes, dehydrogenases*. In: (Miller RH & Keeney DR Eds., *Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties*, Agronomy Monograph, No. 9, ASA and SSSA, Madison), 1982.
- 42 Tabatabai MA & Bremner JM, Assay of urease activity in soils. *Soil Biol Biochem*, 4 (1972) 479.
- 43 Vance EM, Brookes PC & Jenkinson DS, An extraction method for measuring soil microbial biomass C. *Soil Biol Biochem*, 19 (1987) 703.
- 44 Brookes PC, Landman A, Pruden G & Jenkinson DS, Chloroform fumigation and the release of soil nitrogen: a rapid direct extraction method to measure microbial biomass nitrogen in soil. *Soil Biol Biochem*, 17 (1985) 837.
- 45 Spaepen S, Das F, Luyten E, Michiels J & Vanderleyden J, "Indole-3-acetic acid-regulated genes in *Rhizobium etli* CNPAF512: research Letter". *FEMS Microbiology Lett*, 291 (2009) 195.
- 46 Kumar PR & Ram MR, Production of indole acetic acid by *Rhizobium* isolates from *Vigna trilobata* (L.) Verdc. *Afr J Microbiol Res*, 6 (2012) 5536.
- 47 Sarkar D & Laha S, Production of phytohormone auxin (IAA) from soil born *Rhizobium* sp. isolated from different leguminous plant, *Int J Appl Environ Sci*, 8 (2013) 521.
- 48 Ghosh PK, De TK & Maiti T K, Production and metabolism of indole acetic acid in root nodules and symbiont (*Rhizobium undicola*) isolated from root nodule of aquatic medicinal legume *Neptunia oleracea* Lour. *J Bot*, (2015) 1.
- 49 Kucuk C & Cevheri C, Indole acetic acid production by *Rhizobium* sp. isolated from pea (*Pisum sativum* L. ssp. *arvense*). *Turk J Life Sci*, 1(2016) 43.
- 50 Shamseldin A & Werner D, High salt and high pH tolerance of new isolated *Rhizobium etli* strains from Egyptian soils, *Curr Microbiol*, 50 (2005) 11.
- 51 Belal EB, Hassan MM & El-Ramady HR, Phylogenetic and characterization of salt-tolerant rhizobial strain nodulating faba bean plants, *Afr J Biotech*, 12 (2013) 4324.
- 52 Guerrero-Castro J, Lozano L & Sohlenkamp C, Dissecting the acid stress response of *Rhizobium tropici* CIAT 899. *Front Microbiol*, 9 (2018) 1.
- 53 Omotoyinbo OV & Omotoyinbo BI, Effect of Varying NaCl Concentrations on the Growth Curve of *Escherichia coli* and *Staphylococcus aureus*, *Cell Biol*, 4 (2016) 31.
- 54 Rai R, Dash PK, Mohapatra T & Singh A, Phenotypic and molecular characterization of indigenous rhizobia nodulating chick pea in India. *Indian J Exp Biol*, 50 (2012) 340.
- 55 Patil SM, Patil DB, Patil MS, Gaikwad PV, Bhamburdekar SB & Patil PJ, Isolation, characterization and salt tolerance activity of *Rhizobium* sp. from root nodules of some legumes. *Int J Curr Microbiol App Sci*, 3 (2014) 1005.
- 56 Chu HY, Lin XG, Fujii T, Morimoto S, Yagi K, Hua JL & Zhang JB, Soil microbial biomass, dehydrogenase activity, bacterial community structure in response to long-term fertilizer management. *Soil Biol Biochem*, 39 (2007) 2971.
- 57 Yang RY, Tang JJ, Chen X & Hu SJ, Effects of coexisting plant species on soil microbes and soil enzymes in metal lead contaminated soils. *Appl Soil Ecol*, 37 (2007) 240.
- 58 Bastida F, Kandeler E, Moreno JL, Ros M, García C & Hernández T, Application of fresh and composted organic wastes modifies structure, size and activity of soil microbial community under semiarid climate. *Appl Soil Ecol*, 40 (2008) 318.
- 59 Teng Y, Shen Y, Luo Y, Sun X, Sun M, Fu D, Li Z & Christie P, Influence of *Rhizobium meliloti* on phytoremediation of polycyclic aromatic hydrocarbons by alfalfa in an aged contaminated soil. *J Hazard Mat*, 186 (2011) 1271.
- 60 Bakhoun N, Ndoye F, Kane A, Assigbetse K, Fall D, Sylla SN, Noba K & Dioul D, Impact of rhizobial inoculation on *Acacia senegal* (L.) Willd. growth in greenhouse and soil functioning in relation to seed provenance and soil origin. *World J Microbiol Biotechnol*, 28 (2012) 2567.
- 61 Fall D, Bakhoun N, Sall SN, Zoubeirou AM, Sylla SN & Dioul D, Rhizobial inoculation increases the soil microbial functioning and gum arabic production of 13 year old *Senegalia Senegal* (L.) Britton, trees in north part of Senegal. *Front Plant Sci*, 7 (2016) 1365.
- 62 Gopalakrishnan S, Sathya A, Vijayabharathi R, Varshney RK, Laxmipathi GCL & Krishnamurthy L, Plant growth promoting rhizobia: Challenges and opportunities. *3 Biotech*, 5 (2015) 355.