



Stock assessment of Indian squid, *Uroteuthis (Photololigo) duvaucelii* (d'Orbigny [in Férussac & d'Orbigny], 1835) from south-western Bay of Bengal

E M Chhandaprajnadarsini^{*:a}, N Rudramurthy^a, S Sethi^b, S J Kizhakudan^a & M Sivdas

^aICAR-Central Marine Fisheries Research Institute, P. B. No. 1603, Ernakulam North P. O., Cochin, Kerala – 682 018, India

^bICAR-Central Institute of Brackishwater Aquaculture MRC Nagar, Raja Annamalai Puram, Chennai, Tamil Nadu – 600 028, India

*[E-mail: elinamadhu1@gmail.com]

Received 26 July 2019; revised 24 February 2020

Stock assessment of the Indian squid *Uroteuthis (Photololigo) duvaucelii* off north Tamil Nadu coast from commercial trawl landings was studied during the period 2012-2016. The species is mainly caught by trawl net and the annual average landing of the species was 563.3 t contributing to 2 % to the total trawl landing along north Tamil Nadu coast. L_{∞} , K and t_0 were estimated as 260 mm, 0.84 yr^{-1} and -0.105 yr , respectively. Total mortality rate (Z), Natural mortality rate (M), and Fishing mortality rate (F) were 4.43 yr^{-1} , 1.67 yr^{-1} and 2.78 yr^{-1} , respectively. t_{max} was estimated to be 3.47 yr. The length at first capture ($L_{c50} = 62 \text{ mm}$) estimated by Length converted catch curve method was found to be lower than Length at first maturity ($L_{m50} = 80 \text{ mm}$). Furthermore, E_{curr} (current exploitation rate) was found to be on higher side than the E_{max} (0.48) and $E_{0.1}$ (0.40) which indicates that the fishery is in overexploited stage. Thompson and Bell prediction model showed that a marginal decrease (20 %) in current level of exploitation would help in regeneration of stock for long term sustainability of the resources as well as in achieving maximized economic return.

[**Keywords:** Exploitation rate, Spawning stock biomass, *Uroteuthis (Photololigo) duvaucelii*]

Introduction

Overfished stocks of marine fisheries present great concern globally and about 33.1 % of stocks are currently being overfished¹. There has been a steady increase in percentage of stock fished without giving concern to the sustainability. Although, fisheries resources are renewable resources, such huge irremediable depletion of fish stocks as a result of overexploitation has led to decline in fishery resources all over the world. This is where, stock assessment played an important role by giving information regarding status of the fishery *i.e.* whether it is being over exploited or underexploited. It also tells us the possible outcomes of different management actions to the fisheries managers.

Cephalopods belongs to the phylum mollusca and are represented by 800 species comprising of three main categories *i.e.* squid, cuttlefish and octopus. The commercial exploitation of cephalopods has been steadily increasing due to ever increasing demand in internal and external market. With the increasing demand for frozen cephalopods in the export basket to several countries, the fishery has transitioned from a by-catch resource to a targeting resource. Off late cephalopods product has become a major item

exported from India. The resources are intensively exploited by single day and multiday trawlers. Among the cephalopods, cuttlefish formed the major fishery followed by squids and octopus along Chennai coast. *U.(P.) duvaucelii*, the Indian squid, is extensively distributed in neritic waters from Mozambique to the South China Sea and from the Philippines Sea up to Taiwan; occurring up to a depth of 120 m or even more². In Tamil Nadu it is locally known as “Usikaddama” mainly exploited by single day and multiday trawlers. Several authors³⁻⁷ have studied the fishery and population dynamics of Indian squid in Indian waters. However, the information on stock status of *U.(P.) duvaucelii* along the east coast is scarce. Studies on the stock status of commercially important species are essential for the long term sustainability of fishery resources and fishing industry. Furthermore, the paucity of information on population parameters and stock status of the Indian squid especially within east coast of India may cripple any management intervention towards the sustainable exploitation of the resources in India. It is against this backdrop, current study has been conducted to estimate the stock parameters of *U.(P.) duvaucelii* to enhance the existing management intervention.

Materials and Methods

Weekly sampling was carried out from the commercial trawl landing at Madras fisheries Harbour (MFH) in Kasimedu, Chennai for a period of 5 years during January, 2012 to December, 2016 to collect the data on catch, effort and size composition of Indian squid (Fig. 1). Dorsal mantle length and the individual body weight was measured from fresh specimens of *U.(P.) duvaucelii* to the nearest mm and mg, respectively. An eye estimated catch of the species was recorded on the day of observation. Daily length frequency data collected for the species were raised by the raising factor (the total landing weight / sample weight of the species on the day of observation). Afterwards, the raised daily length frequency data were again raised to monthly landing, following the method adopted by Fisheries Resources Assessment Division (FRAD) of CMFRI⁸. The von Bertalanffy growth parameters (L_{∞} and K) were estimated using monthly raised length frequency data in the ELEFAN 1 module of FiSAT II⁹. The age at length zero (t_0) was estimated by using Pauly's empirical equation: $\text{Log}_{10}(-t_0) = -0.3922 - 0.2752 \log_{10} L_{\infty} - 1.038 \log_{10} K^{\text{(ref. 10)}}$. Growth performance index (ϕ) was estimated from the formula: $\phi = \log_{10} K + 2 \log_{10} L_{\infty}^{\text{(ref. 11)}}$. Longevity (t_{max}) was calculated from the formula $t_{\text{max}} = 3/K + t_0^{\text{(ref. 12)}}$. The length-weight relationship of *U. duvaucelii* was established using the formula, $W = aL^{\text{b(ref. 13)}}$.




Fig. 1 — Map showing Kasimedu fishing harbour (13° 07' 32'' N & 80° 17' 49'' E) south western Bay of Bengal

The raised annual length frequencies of the Indian squid were pooled for all the years (2012-2016) and the pooled length frequency values were considered for further study. Instantaneous total mortality rate (Z) and Instantaneous natural mortality rate (M) were estimated by length converted catch curve method¹⁴ and Pauly's empirical equation¹⁵, respectively. The fishing mortality (F) was calculated as $F = Z - M$. Following the method described by Beverton & Holt¹⁶, the exploitation rate ($U = F/Z (1 - e^{-Z})$) and exploitation ratio ($E = F/Z$) were estimated. Using FiSAT, recruitment pattern of Indian squid was determined by backward projection on length axis of length frequency data. The restructured data was used to reduce the temporal spread and recruitment pattern was determined by maximum likelihood method using NORMSEP¹⁷. The length at recruitment (L_r) was considered to be the midpoint of the smallest length group in the catch during the five-year period. Length at first capture (L_{C50}) was estimated from length converted catch curve method¹³. The probability of capture was estimated by backward extrapolation of regression line of descending limb of length converted catch curve and by using logit curve, the probability of capture of sequential length classes were regressed to obtain L_{C50} . Size at first maturity (L_{M50}), defined as the length at which 50 % of the female stock attain sexual maturity, was determined by graphical method¹⁸. The cumulative percentage frequency of II to IV stages were plotted against various length groups and the length at 50 % cumulative percentage frequency was considered to be length at first maturity. Maturity stages of female were assessed by visualising its internal organ and classified into the four point maturity scale following the method described by Silas for squid¹⁹.

Based on the estimated growth parameters and fishery parameters of the species, the relative yield per recruit (Y'/R) and relative Biomass per recruit (B'/R) were estimated at different fishing levels using Beverton & Holt's relative yield per recruit analysis method²⁰. By using Thompson & Bell bio-economic model²¹, Yield (Y), Biomass (B) and spawning stock biomass (SSB) were predicted at different exploitation levels.

Results and Discussion

Fishery

The data on annual landing and annual effort data for total squids as well as *U.(P.) duvaucelii* during

2012-2016 is shown in Table 1. The analysis indicated that annual squid landing at MFH fluctuated between 732 and 1671 t during 2012-2016, with maximum landing in 2015. Along north Tamil Nadu coast cephalopods were mainly targeted by multiday and single day trawl net and *Sepia pharaonis* forms the major fishery followed by *U.(P.) duvaucelii* among cephalopods in north Tamil Nadu coast. The average landing of *U.(P.) duvaucelii* was 1050.6 t during 2012-2016 with an average CPUE of 1.14 kg hr⁻¹ (Table 1). Although maximum landing of the species was recorded in 2015, its percentage contribution to the total squid landing of north Tamil Nadu was lowest in that year, where it formed only about 38 %; while in the other years it formed 53-68 %. It was because of more quantity of other species of squid landed along the coast during 2015. Generally along North Tamil Nadu Indian squid forms the major fishery and it contributes more than 65 % of the landing while other species viz. *U.(P.) edulis*, *U.(P.) singhalensis* and *Loliolus (Nipponololigo) uyii* landed in insignificant quantities.

Length -weight relationship (LWR)

The relationship between dorsal mantle length (ML) and total body weight (W) was estimated for pooled individuals as well as for both individual sexes (male and female) separately. The length-weight relationship was estimated based on 153 males (in the ML range of 13.5 to 26.6 cm and weight in the range of 14 to 117 g) and 319 females (in the ML range of 12.8 to 31 mm and weight in the range of 12 to 214 g).

The LWR in Indian squid can be expressed as:

$$\text{Male: } W = 0.35(\text{ML})^{1.92} \quad (R = 0.87)$$

$$\text{Female: } W = 0.14(\text{ML})^{2.54} \quad (R = 0.91)$$

$$\text{Pooled: } W = 0.19(\text{ML})^{2.19} \quad (R = 0.89)$$

The value of b found in present study for male, female and pooled sexes indicates negative allometric

growth pattern in Indian squid. However male showed higher degree of negative allometric growth pattern compared to the female. The statistical comparisons indicated a significance difference ($p = 0.0009 < 0.01$) between LWR of male and female. The LWR of *Uroteuthis (Photololigo) duvaucelii* was studied by several authors^{4,22-26} and they reported negative allometric growth pattern, as found in the present study.

Growth parameters

A total of 2093 Indian squids were randomly collected for the current study and the specimens which are used for the present study were in the size range of 25-239 mm. The restructured length frequency distribution and growth performance of *U.(P.) duvaucelii* is shown in Figure 2. The von Bertalanffy growth parameters (L_{∞} and K) were estimated at 260 mm and at 0.84 y⁻¹, respectively. The t_0 was estimated as -0.105 yr while longevity (t_{\max}) was estimated to be 3.47 yr. Thus, von Bertalanffy equation for *U.(P.) duvaucelii* from south-western Bay of Bengal can be expressed as $L_{\infty} = 260 * (1 - e^{-(0.84 - (t - (-0.105)))})$. The growth curve indicated that *U.(P.) duvaucelii* grows to a ML of 157.32 mm in 1st year, 215.63 mm in 2nd year and 240.85 mm at the end of 3 years of its life. Thomas & Kizhakudan⁷ reported that the Indian squid grows upto 189, 260, 287 mm at the end of the first three years of its life along the Gujarat coast, with longevity of 3.06 years. However, Karnik *et al.*²⁷ reported the length at the end of I to III years for *U.(P.) duvaucelii* as 221, 315 and 355 mm along Mumbai coast, with longevity of 3.5 years.

The L_{∞} estimated in the current study is comparatively smaller than the earlier reports^(2,3,6,7,22,27-33) except Meiyappan *et al.*⁴ from the east coast (Table 2). They reported L_{∞} of 220 mm for male and 205 mm for female from east coast of India and higher values of 360 and 232 mm for male and female respectively from west coast of India. An earlier study by Mohamed³⁴ indicated that females grew faster than males, while males ultimately reach larger size. As

Table 1 — Catch and effort of *U.(P.) duvaucelii* along south western Bay of Bengal

Year	Effort (fishing hours)	Total trawl landing (t)	Squid landing (t)	<i>U.(P.) duvaucelii</i>		
				Landings (t)	CPUE (kg/hr)	% contribution to total squid landing
2012	422819	23124	732	435	1.03	59.41
2013	470457	26277	989	526	1.12	53.18
2014	432884	28985	940	637	1.47	67.82
2015	476284	29386	1671	639	1.34	38.24
2016	783387	35348	921	579	0.74	62.87
Average	517166.2	28624	1050.6	563.2	1.14	56.30




Fig. 2 — Restructured length frequency of *U.(P.) duvaucelli*

Table 2 — A summary of earlier reports on growth and mortality parameters

Authors	Region	Sex	L_{∞} (mm)	K (yr ⁻¹)	Z (yr ⁻¹)	M (yr ⁻¹)	F (yr ⁻¹)	t_0 (yr)	t_{max} (yr)
Silas <i>et al.</i> ²²	Cochin coast	M	327	0.61					
		F	205	1.19					
Kasim ³	Veraval coast	Pooled	300	0.98	3.94	1.74	2.20	0.56	3.10
Supongpan ²⁸	Thailand coast	Pooled	266	0.86					3.49
Meiyappan & Srinath ²	Cochin coast	M	379	1.10	9.0				
		F	238	1.17	10.6				
Vidyasagar & Deshmukh ²⁹	Mumbai coast	Pooled	323	0.45	1.83				
Meiyappan <i>et al.</i> ⁴	East coast of India	M	220	0.90					
		F	205	1.30					
Chakraborty <i>et al.</i> ³⁰	Mumbai coast	Pooled	343	0.49	2.90	1.10	0.99		
Mohamed & Rao ⁶	Karnataka coast	Pooled	343	1.43					2.50
Karnik <i>et al.</i> ²⁷	Mumbai coast	Pooled	385	0.85	4.29	1.82	2.47		3.53
Thomas & Kizhakudan ⁷	Veraval coast	Pooled	303	0.98	3.94	1.74	2.20		3.06
John ³¹	Cochin	Pooled		0.75	2.61	1.37	1.24		
Chakraborty <i>et al.</i> ³²	Mumbai	Pooled	380	0.90	4.43	1.41	3.02		3.33
Nitin <i>et al.</i> ³³	India	Pooled	376	0.95	4.50	1.82	2.68		3.10
Present study	North Tamil Nadu, South western Bay of Bengal	Pooled	260	0.84	4.43	1.67	2.76	-0.105	3.47

the estimation of L_{∞} is greatly influenced by L_{max} in the sample, L_{∞} may vary from location to location.

Similarly the value of growth coefficient ‘K’ obtained in this study is found to be on higher side compared to the value estimated by Vidyasagar & Deshmukh²⁹ and Chakraborty *et al.*³⁰ from Mumbai waters and by John³¹ from Cochin waters. However, Mohamed & Rao⁶ estimated significantly higher value of K from Karnataka coast. The estimated value of K in the current study is in agreement with Karnik *et al.*²⁷ and Supongpan²⁸. Growth parameters may vary for different stocks belonging to different geographic locations. It has been reported that in

some squids, particularly those belonging to family Loliginidae, the growth parameters are highly variable for different stocks³⁵.

Mortality parameters

By using length converted catch curve¹⁴ method, the total mortality rate was estimated to be 4.43 (Fig. 3). The natural mortality (M) and fishing mortality (F) were estimated as 1.65 and 2.78, respectively. The present obtained values of natural mortality, fishing mortality and total mortality has been compared with earlier studies (Table 2). The total annual mortality of the Indian squid obtained in the current study is comparatively on higher side than the




Fig. 3 — Estimation of Z from length converted catch curve ($Z = 4.43$)

earlier reports^(2,3,7,29-31). Nevertheless, the present estimated value of total mortality is slightly closer to the earlier reports by Chakraborty *et al.*³² and Karnik *et al.*²⁷. Natural mortality was compared with earlier studies and it seems to be different for various geographic locations. The variation in natural mortality observed in the present study may be because of environmental conditions as well as fishing methods which influence the abundance of predators and competitors³⁶. *U.(P.) duvaucelii* is a preferred food of most fishes in the marine ecosystem, and the abundance of predators may affect the natural mortality. Nevertheless, M/K is used as an index of reliable estimation of M and K value and it ranges from 1 to 2.5^(ref. 36). The present estimated value of M/K (1.96) indicated a reasonable estimation of natural mortality.

The current fishing mortality rate ($F = 2.78$) is comparatively higher than natural mortality ($M = 1.63$) revealing overexploitation of Indian squid. However, a stock is considered to be in a healthy state and at optimum exploited phase, if the fishing mortality and natural mortality are equal in a natural ecosystem (*i.e.* $E = 0.5$)³⁷. However, the present value of exploitation rate is comparatively higher than the optimum exploitation rate affirming the evidence of overfishing of squid along the north Tamil Nadu coast.

Length at first capture (L_{c50}) and Length at first maturity (L_{m50})

The length at first capture *i.e.* length at which 50 % of the squid in the stock become vulnerable to the gear was found to be 62 mm while maturity studies indicated that 50 % of the squid attain maturity at

DML of 80 mm. The L_{m50} was relatively higher than L_{c50} , indicating that the squid are mostly exploited before they attain maturity, which is clear sign of growth overfishing. Moreover the value of L_c/L_∞ (0.24) was far below the recommended value of 0.5^(ref. 38) indicating that majority of the catch consists of juveniles, a common feature of growth overfishing. The bulk landing of juvenile squid may be due to the use of trawl gear with small mesh size (< 30 mm) which is used to target shrimp, and also due to the use of non-selective gears for squids along the north Tamil Nadu coast. The optimum L_{c50} which corresponds to the mantle length for obtaining maximum possible yield, was estimated to be 157.88 mm. L_{m50} , estimated in the present study is observed to be close to earlier reports^{39,40}. However, Rao⁴¹ reported a higher value of L_{m50} (108 mm) from Mangalore coast. Smaller size at maturity may be due to the effect of high fishing pressure.

Recruitment pattern

The recruitment pattern of *U.(P.) duvaucelii* was continuous throughout the year with, one major peak during June to August and with one minor peak during February to March (Fig. 4). Highest recruitment was found to be in the month of April (14.65 %) and May (23.35 %). The presence of recruits in fishery throughout the year revealed that the Indian squid spawns throughout the year. Silas *et al.*⁴² observed mature gonads in squids almost throughout the year all along the coast which is in agreement with the present study. The observed smallest mantle length at first recruitment (L_r) was 35 mm, providing the evidence of the use of small mesh trawl gears by fishermen along the north Tamil Nadu coast.

Stock assessment

By using input parameters of L_c/L_∞ (0.24) and M/K (1.96) the Beverton and Holt's relative yield per recruit and biomass per recruit analysis was carried out for the stock assessment of *U.(P.) duvaucelii*. It showed that maximum yield/recruit could be obtained at exploitation rate of 0.48; while the optimum yield *i.e.*, half of its virgin biomass, can be obtained at exploitation rate of 0.29 (Fig. 5). Nevertheless, as the aquatic ecosystem is highly uncertain, as precautionary approach $E_{0.1} = 0.40$ can be taken as management reference point. The current exploitation rate ($E = 0.63$) is relatively higher than this reference point, indicating the overexploited status of the Indian squid in the south-west Bay of Bengal along the north Tamil Nadu coast.




Fig.4 — Recruitment curve of *U.(P.) duvaelii* along south western Bay of Bengal coast




Fig. 5 — Yield per recruit and biomass per recruit showing the maximum yield per recruit at exploitation rate levels $\{E_{max} = 0.48, E_{10} = 0.40, E_{50} = 0.29\}$ of *U.(P.) duvaelii* from the North Tamil Nadu, south west Bay of Bengal

Managing fisheries under uncertainty has become a top priority and a major challenge in recent times. The Beverton and Holt’s relative Y/R and B/R analysis has some limitation while estimating management reference points as it tends to overestimate the E_{max} unrealistically for smaller species with high natural mortality in tropical ecosystem. Moreover, the above analysis does not consider spawning stock biomass⁴³⁻⁴⁴; whereas spawning stock biomass is an important indication of health status of the stock. There should be sufficient spawning stock biomass in the ecosystem for the regeneration of exploited stock and it is always advisable to maintain the virgin spawning stock




Fig. 6 — Length based Thompson and Bell analysis of *U.(P.) duvaelii* along North Tamil Nadu, south western Bay of Bengal coast predicting the yield, spawning stock biomass and total stock biomass at different effort level




Fig. 7 — Percentage of initial biomass spawning stock biomass of *U.(P.) duvaelii* at different fishing effort level of F

biomass at 20-30 % in the ecosystem to minimize the risk of recruitment overfishing and to ensure the availability of adequate amount of spawners in the ecosystem⁴⁵.

Due to risk uncertainty in fisheries stock assessment, Thompson & Bell bio-economic model²¹ was used to study the impact of different fishing levels on Yield (Y), Biomass (B) and spawning stock biomass (SSB) and the results (Fig. 6) indicated that the current fishing effort can be increased by 20 % to obtain Maximum Sustainable Yield. However, this may lead to a decline in the spawning stock biomass to 13.4 %, which may be detrimental to the prolong sustainability of Indian squid fishery along the coast. In order to achieve Maximum Economic Yield, the current fishing level needs to be decreased by 20 % and at that level spawning stock biomass could be maintained at 23.78 % (Fig. 7). The marginal decrease

in effort by 20 % would help in regeneration of stock as well as in achieving maximized economic return. However, in a tropical country like India where multispecies multi-gear fisheries exist, stock status of other resources inhabiting the same ecosystem should be carefully studied before arriving at any management intervention, as the decrease or increase in effort level may affect the fishery of other species.

Conclusion

Sustainability is a major goal of fisheries management. The present study indicates that the continued use of trawl gear with small mesh size for fishing as well as higher fishing effort can lead to lower economic return, reduction in CPUE (Catch per unit effort) and can eventually lead to a collapse of fishery of Indian squid in the near future. Therefore it is advisable to reduce the effort level in order to minimise the risk of growth overfishing as well as recruitment overfishing, since large mesh fishing gear targeting large sized individuals will allow the juvenile fish to grow and spawn at least once, ensuring sustainability of the resource.

Acknowledgments

The authors are grateful to Director, ICAR-CMFRI, Cochin for providing constant support and all necessary facilities during the entire study.

Conflict of Interest

There is no conflict of interest.

Author Contributions

Data collection: EMC, NR and SS. EMC: Data analysis and wrote the paper. SJK and MS: reviewed the manuscript and coordinated the work.

References

- 1 FAO, *The State of World Fisheries and Aquaculture-Meeting the sustainable development goals*, (Food and Agriculture Organisation, Rome), 2018, pp. 210.
- 2 Meiyappan M M & Srinath M, Growth and mortality of the Indian squid (*Loligo duvaucelii*) off Cochin, India, In: *Contributions to tropical fish stock assessment in India: FAO/DANIDA/ICAR National Follow-up Training Course on Fish Stock Assessment*, (1989), pp. 1-14.
- 3 Kasim H M, Population dynamics of the squid *Loligo duvaucelii* d'Orbigny (Cephalopoda) in Saurashtra waters, *J Mar Biol Assoc India*, 2 (1) (1985) 103-112.
- 4 Meiyappan M M, Srinath M, Nair K P, Rao K S, Sarvesan R, *et al.*, Stock assessment of the Indian squid *Loligo duvaucelii* d'Orbigny, *Indian J Fish*, 4 (1) (1993) 319-329.
- 5 Mohamed K S, Estimates of growth, mortality and stock of the Indian squid *Loligo duvaucelii* d'Orbigny, exploited off Mangalore, southwest coast of India, *Bull Mar Sci*, 2 (1) (1996) 393-403.
- 6 Mohamed K S & Syda G R, Seasonal growth, stock-recruitment relationship and predictive yield of the Indian squid *Loligo duvaucelii* d'Orbigny exploited off Karnataka coast, *Indian J Fish*, 4 (1) (1997) 319-329.
- 7 Thomas S & Shoba J K, Cephalopod fishery and population dynamics of *Loligo duvaucelii* (d'Orbigny) off Saurashtra region, Gujarat, *Indian J Fish*, 4 (1) (2006) 425-430.
- 8 Sekharan K V, On oil sardine fishery of the Calicut area during the year 1955-56 to 1958-59, *Indian J Fish*, 9 (2) (1962) 679-700.
- 9 Gayanilo F C, Per S & Daniel P, *FAO-ICLARM stock assessment tools II: User's guide*, (Food & Agriculture Organisation, Rome), 2005, pp.168.
- 10 Pauly D, A selection of simple methods for the assessment of tropical fish stock, *FAO Fisheries Circular*, 729 (1980) 54.
- 11 Pauly D & Munro J L, Once more on the comparison of growth in fish and invertebrates, *Fishbyte*, 1 (1984) 1-21.
- 12 Pauly D, *Some simple methods for the assessment of tropical fish stocks*, (Food & Agriculture Organisation, Rome), 1983, pp. 11.
- 13 Le Cren E D, The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*), *J Anim Ecol*, 20 (1951) 201-219.
- 14 Pauly D, Length-converted catch curves: a powerful tool for fisheries research in the tropics, *Fishbyte*, 2 (1) (1983) 9-13.
- 15 Pauly D, On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *ICES J Mar Sci*, 2 (1980) 175-192.
- 16 Beverton R J & Sidney J H, *On the dynamics of exploited fish populations*, (Springer Science & Business Media, London), 1957, pp. 533.
- 17 Pauly D & Caddy J F, *A modification of Bhattacharya's method for the analysis of mixtures of normal distributions*, (Food and Agricultural Organization, Rome), 1985, pp. 16.
- 18 Udupe K S, Statistical method of estimating the size at first maturity in fishes, *Fishbyte*, 2 (1986) 8-10.
- 19 Silas E G, Cephalopod fisheries of India - An introduction to the subject with methodologies adopted for the study. In: *Cephalopod bionomics, fisheries and resources of the EEZ of India*, edited by E G Silas, *Bull Cen Mar Fis Res*, 37 (1985) 1-4.
- 20 Beverton R J H & Holt S J, *Manuals of methods for fish stock assessment: Part II Tables of yield function*, *FAO Fish Biol Tech Paper*, 38 (1966) 67.
- 21 Thompson W F, Biological statistics of the pacific halibut fishery: Effect of changes in intensity upon total yield and yield per unit of gear, *Rep Int Fish Commn*, 8 (1934) 1-49.
- 22 Silas E G, Rao K S, Sarvesan R, Nair K P, Vidyasagar K, Meiyappan, *et al.*, Some aspects of the biology of squids, *CMFRI Bull*, 37 (1985) 38-48.
- 23 Nair K, Meiyappan M M, Kuriakose P S, Sarvesan R, Lipton A P, *et al.*, Biology of squids. Exploratory squid jigging in India with notes on biology of squids, *Fisheries Survey of India*, 23 (1992) 27-42.
- 24 Srichanngam S, *Age and growth determination and stock identification using statolith microstructure of Indian squid, Loligo duvaucelii*, MS thesis, The University of Bergen, Norway, 2010.
- 25 Mishra A S, Prakash N & Somvanshi V S, Length-weight relationship, condition factor and sex ratio of *Uroteuthis (Photololigo) duvaucelii* (d'Orbigny, 1848) from Goa, west coast of India, *Mar Boil Assoc India*, 2 (2012) 65-68.

- 26 Petsut N & Kulabtong S, Biology of Indian squid, *Loligo duvaucelii* in Thailand, *Veridian*, 2 (2012) 1005-1009.
- 27 Karnik N S, Chakraborty S K, Jaiswar A K, Swamy R P, Rajaprasad R, *et al.*, Growth and mortality of Indian squid, *Loligo duvaucelii* (d'Orbigny) (Mollusca/ Cephalopoda/ Teuthoidea) from Mumbai waters, India, *Indian J Geo-Mar Sci*, 1 (2003) 67-70.
- 28 Supongpan M, Assessment of Indian squid (*Loligo duvaucelii*) and Mitre squid (*L. chinensis*) in the Gulf of Thailand, In: *FAO Fisheries Report*, edited by S Venema, J Moller Christensen & D Pauly, 1988, pp. 25.
- 29 Vidyasagar K & Deshmukh V D, Stock assessment of *Loligo duvaucelii* (d'Orbigny) in Bombay waters, *J Mar Biol Assoc India*, 2 (1992) 14-17.
- 30 Chakraborty S K, Deshmukh V D, Khan M Z, Vidyasagar K & Raje S G, Estimates of growth, mortality, recruitment pattern and maximum sustainable yield of important fishery resources of Maharashtra coast, *Indian J Geo-Mar Sci*, 26 (1997) 53-56.
- 31 John M, *Studies on Some Aspects of Landings, Utilization and Export of Commercially Important Cephalopods*, Ph.D. thesis, Cochin University of Science and Technology, Kerala, India, 2007.
- 32 Chakraborty S K, Biradar R S, Jaiswar A K, Palaniswamy R & Pawan K, Growth, mortality and population dynamics of three cephalopods species, *Loligo duvaucelii* (Orbigny), *Sepia aculeate* (Orbigny) and *Sepiella innermis* (Orbigny) from north- west coast of India, *Indian J Fish*, 3 (2013) 1-7.
- 33 Nitin P, Nirmale V H, Metar S Y, Bhsale B P, Sawant M S, *et al.*, Age, growth and mortality studies of Indian squid, *Uroteuthis (Photololigo) duvaucelii* (d'Orbigny) along Ratnagiri Coast of Maharashtra, India, *Indian J Geo-Mar Sci*, 1 (2015) 93-96.
- 34 Mohamed K S, Estimates of growth, mortality and stock of Indian squid *Loligo duvaucelii* (Orbigny), exploited off Mangalore. South west coast of India, *Bull Mar Sci*, 2 (1996) 393-403.
- 35 Pauly D, Population dynamics of short-lived species, with emphasis on squids, *J Northwest Atl Fish Sci*, 106 (1) (1984) 101-106.
- 36 Beverton R J H & Holt S J, *A review of the lifespan and mortality rates of fish in nature, and their relation to growth and other physiological characteristics*, (CIBA Foundation colloquia on ageing), 5 (1959) pp. 142-177.
- 37 Beverton R J H & Holt S J, On the dynamics of exploited fish populations, *Fish Invest Ser Mar Fish GB Minist Agric Fish*, 19 (1957) 533.
- 38 Baijot E & Moreau J, Biology and demographic status of the main fish species in the reservoirs of Burkina Faso, (Hydrobiological aspects of fisheries in small reservoirs in the Sahel region. Technical Center for Agricultural and Rural Cooperation ACP-EU, Wageningen, Netherlands), 1997, pp. 79-110.
- 39 Silas E G, Rao S, Servesan K, Prabhakaran R, Nair K, *et al.*, The exploited squid and cuttlefish resources of India - A review, *Mar Fish Infor Sew T & E Ser*, (1982) 1-16.
- 40 Naik N R, Shivaprakash S M, Anjaneyappa H N, Somasekhara, S R, Naik J, *et al.*, Reproductive biology of the commercially important Indian squid *Uroteuthis (Photololigo) duvaucelii* off Mangalore, south-west coast of India, *Indian J Fish*, 64 (2017) 75-79.
- 41 Rao G S, Biology of inshore squid *Loligo duvaucelii* Orbigny, with a note on its fishery off Mangalore, *Indian J Fish*, 3 (1988) 121-30.
- 42 Silas E G, Rao S, Servesan K, Prabhakaran R, Nair K, *et al.*, Some aspects of the biology of squids. In: *Cephalopods-Bionomics, fisheries and resources of the exclusive economic zone of India*, edited by E G Silas, *CMFRI Bulletin*, 37 (1985) 38-48.
- 43 Gulland J A, *Fish stock assessment: a manual of basic methods*, (FAO Wiley series on food and agriculture), 1983, pp. 223.
- 44 Pauly D, *Fish population dynamics in tropical waters: a manual for use with programmable calculators*, *ICLARM Stud Rev*, 8 (1984) 325.
- 45 Rosenberg A A & Victor R, Precautionary management reference points and management strategies, *FAO Fish Tech Pap*, 350 (1996) 129-140.