

Electronic Supplementary Data

Synthesis and spectroscopic investigation of binding of novel Thiazolo [2, 3-a] isoquinoline analog with bovine serum albumin

Prasanna B Ranade^a, Dinesh N Navale^a, Santosh W Zote^b, Dnyaneshwar K Kulal^c, Mohmad V Sheikh^c & M M V Ramana^{d,*}

^aDepartment of Chemistry, Vivekanand Education Society's College of Arts, Science and Commerce, Chembur, Mumbai 400 071, India

^bDepartment of Chemistry, Sathaye College Vile Parle (East), Mumbai 400 057, India

^cDepartment of Chemistry, Ramnarian Ruia Autonomous College, Matunga (East) Mumbai 400 019, India

^dDepartment of Chemistry, University of Mumbai, Vidyanagari, Santacruz (East), Mumbai 400 098, India

*E-mail: prasannaranade@yahoo.com

S. No.	Contents	Pg. No.
1	Fig. S1 — IR spectra of 10b-cyclobutyl-5, 6-dihydro-2H-thiazolo [2, 3-a] isoquinolin-3(10bH)-one	2
2	Fig. S2 — ¹ H NMR spectra of 10b-cyclobutyl-5, 6-dihydro-2H-thiazolo [2, 3-a] isoquinolin-3(10bH)-one	2
3	Fig. S3 — ¹³ C NMR spectra of 10b-cyclobutyl-5, 6-dihydro-2H-thiazolo [2, 3-a] isoquinolin-3(10bH)-one	3
4	Fig. S4 — GC MS spectra of 10b-cyclobutyl-5, 6-dihydro-2H-thiazolo [2, 3-a] isoquinolin-3(10bH)-one	3




Fig. S1 — IR spectra of 10b-cyclobutyl-5,6-dihydro-2H-thiazolo [2,3-a] isoquinolin-3(10bH)-one




Fig. S2 — ¹H NMR spectra of 10b-cyclobutyl-5,6-dihydro-2H-thiazolo [2,3-a] isoquinolin-3(10bH)-one




Fig. S3 — ^{13}C NMR spectra of 10b-cyclobutyl-5,6-dihydro-2H-thiazolo [2,3-a] isoquinolin-3(10bH)-one




Fig. S4 — GC MS spectra of 10b-cyclobutyl-5,6-dihydro-2H-thiazolo [2,3-a] isoquinolin-3(10bH)-one