

Establishment of an embryogenic suspension culture of *Pinus kesiya* (Khasi pine) from various explants

Chitta Ranjan Deb^{1*} and Pramod Tandon²

¹Botany Department, Nagaland University, Lumami, Mokokchung 798 601, India

²Plant Biotechnology Laboratory, Botany Department, North-Eastern Hill University, Shillong 793 022, India

Received 11 March 2002; accepted 17 June 2003

The embryogenic cultures were obtained from mature zygotic embryos (79.6%) and apical dome sections (92.6%) (pretreated with 0.4% activated charcoal at 4°C for 24 hrs) on mMS and ½DCR media, respectively containing 5 mg l⁻¹ 2,4-D, NAA in combination along with 1 mg l⁻¹ BAP, while, from secondary needles (88.6%) it was on MS medium containing 3 mg l⁻¹ each of 2,4-D and NAA in conjunction with 1 mg l⁻¹ BAP. Using 150-200 mg embryogenic tissues per 15 ml medium at 120 rpm raised embryogenic suspension cultures. But the embryogenic cultures from apical dome sections were cold treated at 4°C for 24 hrs before transferring to suspension culture. The cultures were subcultured at 6-7 days interval and were diluted at 1:4 ratio. The embryonal suspensor masses developed from the resulting elongated single cells within 3-4 passages in respective basal medium with 1/10th growth regulators of initiation medium. The proembryonal head formed in growth regulators free medium at 100 rpm. The embryonal head formed in basal medium containing 4% sucrose and 4 mg l⁻¹ ABA in combination.

Keywords: activated charcoal pre-treatment, cold-treatment, embryogenic suspension culture, *pinus kesiya*, somatic embryogenesis

IPC Code:Int. Cl.⁷ A 01 H 4/00, 7/00

Introduction

Somatic embryogenesis in conifers was first documented from immature zygotic embryos of *Picea abies*¹⁻³. Currently there are about 100 million hectares of industrial tree plantations worldwide, of which 70-80% comprises conifers⁴. For mass clonal propagation of conifers somatic embryogenesis can play a vital role. Since 1985 several papers have been published on successful regeneration of conifer plantlets via somatic embryogenesis both on semi-solid medium and in suspension cultures⁵⁻²⁰.

Pinus kesiya Royle ex Gord. is an economically important early successional tree species, which is predominant in the subtropics (800-2000 m altitude) of North-East India, Myanmar and Philippines. It is the major source of timber in Khasi hills of Meghalaya (India) where it covers about 30% of the total forest area. Some genotypes have tremendous biomass potential and oleo-resin prospects. The Pine forest of the region is getting denudated due to 'Slash and burn' cultivation, ruthless commercial

exploitation and unplanned developmental activities. Therefore, there is an urgent need for large-scale afforestation programme using cloning material. In this paper the authors report establishment of embryogenic suspension culture from zygotic embryos, secondary needles and apical dome sections from mature 'plus' trees of *P. kesiya* (identified by Department of Forestry, Government of Meghalaya).

Materials and Methods

Collection and Priming of Explants

The mature seeds were collected during January-February from mature cones and stored at 4°C in sealed polythene bags till used. The secondary needles were harvested from 5-6-week-old seedlings, during April-May, while 2-3 cm apical domes were collected from 'plus' trees prior to emergence of needles during May-July of every year. Both secondary needles and apical domes immediately after harvesting were placed in anti-oxidant solution (200 mg l⁻¹ PVP) to prevent exudation of phenolic compounds.

Initiation of Culture on Semisolid Phase

The sterilized seeds were imbibed for 24 hrs at 4°C while, the apical domes were cut into 0.5 mm thick

*Author for correspondence:

Tel: 91-369-268221; Fax: 91-369-268204

Email: debchitta@rediffmail.com, dedeb@sancgarbet.in

transverse sections and pretreated on $\frac{1}{2}$ DCR medium²¹ containing 2% sucrose, 0.4% activated charcoal (AC) at 4°C for 72 hrs in the dark before culturing them on growth regulators rich medium for priming the explants. Mature zygotic embryos from imbibed seeds, secondary needles and pretreated apical dome sections were cultured on modified MS [half strength MS salts²² containing 550 mg l⁻¹ NH₄NO₃ and 4460 mg l⁻¹ KNO₃ (mMS)], MS and $\frac{1}{2}$ DCR media, respectively containing 0-3% sucrose, 500 mg l⁻¹ L-glutamine, 1000 mg l⁻¹ casein-hydrolysate and myo-inositol, 0-10 mg l⁻¹ 2,4-D and NAA singly or in combination and 0-5 mg l⁻¹ BAP. Difco-bacto agar (0.7%) was used as gelling agent before autoclaving. Chemicals like ABA, L-glutamine and myo-inositol were filter sterilized and incorporated in the media. In each tube (size-25 × 100 mm) 15 ml medium was poured and for each treatment 50 explants were cultured. The cultures were maintained in the dark at 25±2°C and experiments were repeated thrice. The resulting embryogenic cultures were subcultured biweekly on optimum growth condition for 2 passages. Thereafter, the embryogenic cultures from apical dome sections were cold-treated at 4°C for 0-24 hrs.

Initiation of Suspension Culture

About 50-400 mg embryogenic cultures from above culture conditions were pressed gently with a scalpel blade and transferred to 50 ml conical flask containing 15 ml of respective initiation medium and shaken at 120 rpm on an 'Orbital shaker' (B Braun, model-Certomat U) in the dark at 25±2°C unless mentioned otherwise. The cultures were subcultured at 6-7 day interval for 3-4 passages and were diluted at 1:4 ratio at every subculture.

Maintenance of Cultures

The cultures with embryonal suspension masses (ESMs) elongated single cells from above culture conditions were transferred to respective basal medium containing reduced growth regulators (0, 1/5th and 1/10th of initiation medium) for 4-5 passages. Thereafter, the cultures were grown in growth regulator free medium for 3-4 passages at 100 rpm.

Somatic Embryo Development

The cultures with ESMs were transferred in respective basal medium supplemented with 0-5% sucrose, 0-8 mg l⁻¹ ABA singly or in combination for 6-7 passages.

Results and Discussion

Within two weeks of culture two different types of calli developed. Zygotic embryos and secondary needles formed (i) white, soft and translucent embryogenic (79.6 and 88.6%, respectively) (Figs 1a and b) and (ii) hard and green non-embryogenic cultures. While, from apical dome sections pretreated




Fig. 1—Embryogenic calli formed from different explants on initiation medium, a. from zygotic embryo, b. secondary needles and c. apical dome sections

at 4°C with 0.4% AC formed (i) white, soft and translucent embryogenic (92.6%) (Fig. 1c) and (ii) white but hard non-embryogenic cultures (data not presented). The better embryogenic culture formation from zygotic embryo and apical dome sections were obtained with mMS and ½DCR media, respectively containing 5 mg l⁻¹ each of 2,4-D and NAA along with 2.5 mg l⁻¹ BAP while from secondary needles with MS medium containing 3.0 mg l⁻¹ each of 2,4-D and NAA along with 1.0 mg l⁻¹ BAP. It was observed that apical dome sections primed on activated charcoal rich medium resulted in softer calli when cultured on growth regulators rich medium. It is believed that, during AC pre-treatment the phenolic compounds, growth regulators and other inhibitors adsorbed into the AC, which resulted into soft callus. The promotary effect of AC on the initiation of polyembryogenesis has been documented. Webb and Flinn added 1% AC in the initiation medium which triggered embryogenic culture formation in *P. strobus*²³ while, Gupta *et al* used AC in embryogenic initiation medium for absorbing endogenous ABA and other inhibitory metabolic by-products⁹.

The embryogenic cultures from apical dome sections required a cold treatment at 4°C for 24 hrs before transfer to liquid medium. In the present study, for initiation of suspension cultures, lower inoculum size (150-200 mg/15 ml medium) was found to be suitable at 120 rpm. By decreasing the inoculum (<150 mg/15 ml medium) a poor culture growth was recorded, which may be due to inadequate culture density. Higher inoculum at the initial stage slowed the proliferation of cultures, which may be due to over-crowding of cells. Finer *et al* reported that embryogenic cultures of *P. strobus* proliferated well with low culture density²⁴ while, Krogstrup reported that culture density was crucial and determined the quality of early stage embryos in suspension cultures in *P. sitchensis*²⁵. In the present investigation, the cultures were proliferated by subculturing at 6-7 days interval for 3-4 passages and cultures were diluted at 1:4 ratio at every subculture. During this period cultures yielded elongated embryogenic cells which in turn started cleaving (Fig. 2a) after 2-3 passages in culture. It was necessary to free the single cells from mother tissues at the early stage failing which degeneration of cultures resulted. The ESMs (Figs 2b & c) formed from cleaving cells when they were maintained on respective basal medium with 1/10th growth regulators to that of initiation medium and




Fig. 2—Different stages involved in somatic embryogenesis of *P. kesiya* in suspension culture; a, elongated single cell started cleaving; b and c, embryonal suspensor masses; d, embryonal cell masses showing accumulation; e, somatic embryo with proembryonal head & f, somatic embryo with embryonal head.

maintained for 3-4 passages. The ESMs started accumulation (Fig. 2d) in the medium free of growth regulators and at low agitation where cultures formed proembryonal head. The distinct embryonal head formed (Figs 2e & f) after 5-6 passages when cultures were maintained in respective basal medium containing 4% sucrose and 4 mg l⁻¹ ABA in combination. ABA has been used for cotyledonary embryo maturation in many plant species²⁶ including conifers^{18,27}. In the present study, it was observed that neither sucrose nor ABA singly could promote embryonal head formation but 4% sucrose and 4 mg l⁻¹ ABA in combination was found effective. Tremblay and Tremblay reported that in Black spruce 4-6% sucrose was very effective for maturation of somatic embryos²⁸. Carrier *et al* studied the effect of exogenous sucrose on maturation and germination of somatic embryo of interior spruce and reported that embryos placed on the medium with added sucrose developed roots and epicotyls and increased their

fresh biomass by about 13-fold²⁹. Vagner *et al* reported that with increase in exogenous ABA levels (5-40 μ M), an increase in somatic embryo maturation in *P. abies* both in semisolid and liquid media. In the absence of the exogenous ABA, practically no embryos were formed¹⁶.

In the present study, the initiation, development and maturation of *P. kesiya* somatic embryos have been achieved. This opens up a great potential for large-scale propagation of this species, which is facing rapid denudation.

Acknowledgement

The study was supported by a research grant F. No. BT/R&D/08/003/95 to Prof. Pramod Tandon from Department of Biotechnology, Ministry of Science and Technology, Government of India.

References

- Chalupa V, Somatic embryogenesis and plantlet regeneration from cultured immature and mature embryos of *Picea abies* (L.) Krast, *Communi Inst For Cech*, 14 (1985) 57-63.
- Hakman I, Fowke L C, Von Arnold S & Erikson T, The development of somatic embryos in tissue cultures initiated from immature embryos of *Picea abies* (Norway Spruce), *Plant Sci*, 38 (1985) 53-59.
- Nagmani R & Bonga J M, Embryogenesis in subcultured callus of *Larix deciduas*, *Can J For Res*, 15 (1985) 1088-1091.
- Haines R J, Biotechnology and the genetic improvement of forest trees, *Biotech and Dev Monitor*, 15 (1993) 3-5.
- Attree S M & Fowke L C, Embryogeny of gymnosperms: Advances in synthetic seed technology of conifers, *Plant Cell Tissue Organ Cult*, 35 (1993) 1-35.
- Nagmani R, Diner A M & Sharma G C, Somatic embryogenesis in long leaf pine (*Pinus palustris*), *Can J For Res*, 23 (1993) 873-876.
- Gupta P K & Grob J A, Somatic embryogenesis in conifers, in *Somatic embryogenesis in woody plants*, vol 3 edited by V Jain, P K Gupta & R Newton (Kluwer Academic Publishers, Dordrecht) 1995, 81-98.
- Gupta P K, Timmis R, Timmis K A, Grob J A, Carlson W C *et al*, Clonal propagation of conifers via somatic embryogenesis, in *Proc IUFRO meeting on somatic cell genetics and molecular genetics of trees* (Gent, Belgium) 1995, 55-63.
- Gupta P K, Timmis R, Timmis K A, Carlson W C & Welty E D E, Somatic embryogenesis in Douglas-fir (*Pseudotsuga menziesii*), in *Somatic embryogenesis in woody plants*, vol 3 edited by V Jain, P K Gupta & R Newton (Kluwer Academic Publishers, Dordrecht) 1995, 303-313.
- Keinonen M K, Jalonen P, Eurola P, Von Arnold S & Von Weissenberg K, Somatic embryogenesis of *Pinus sylvestris*, *Scand J For Res*, 11 (1996) 242-250.
- Barrett J D, Park Y S & Bonga J M, The effectiveness of various nitrogen sources in white spruce [*Picea glauca* (Moench) Voss] somatic embryogenesis, *Plant Cell Rep*, 16 (1997) 411-415.
- Guevin T G & Kirby E G, Induction of embryogenesis in cultured mature zygotic embryos of *Abies fraseri* (Pursh) Poir, *Plant Cell Tissue Organ Cult*, 49 (1997) 219-222.
- Norgard J V, Somatic embryo maturation and plant regeneration in *Abies nordmanniana* Lk, *Plant Sci*, 124 (1997) 211-221.
- Li X Y, Huang F H & Gbur J R, Effect of basal medium, growth regulators and phytigel concentration on initiation of embryogenic cultures from immature zygotic embryos of loblolly pine (*Pinus taeda* L.), *Plant Cell Rep*, 17 (1998) 298-301.
- Tandon P & Deb C R, Somatic embryogenesis and plantlet regeneration in *Pinus kesiya* (Royle ex. Gord.), in *Intl Congr on Plant Tissue Cell Cult*, held on 14-19 June, 1998 (Jerusalem) 1998.
- Vagner M, Vondrakova Z, Strnadova Z, Eder J & Machackova I, Endogenous levels of plant growth hormones during early stages of somatic embryogenesis of *Picea abies*, *Adv Hort Sci*, 12 (1998) 11-18.
- Kim Y W, Youn Y, Noh E R & Kim J C, Somatic embryogenesis and plant regeneration from immature zygotic embryos of Japanese larch (*Larix leptolepis*), *Plant Cell Tissue Organ Cult*, 55 (1999) 95-101.
- Arya S, Kalia R K & Arya I D, Induction of somatic embryogenesis in *Pinus roxburghii* Sarg, *Plant Cell Rep*, 19 (2000) 775-780.
- Deb C R & Tandon P, Induction of somatic embryogenesis in Khasi pine (*Pinus kesiya*) from secondary needles, *J Plant Biol*, 29 (2002) 113-118.
- Deb C R & Tandon P, Somatic embryogenesis and plantlet regeneration from mature zygotic embryos of *Pinus kesiya* (Royle ex Gord.), *J Plant Biol*, 29 (2002) 301-306.
- Gupta P K & Durzan D J, Shoot multiplication from mature trees of Douglas-fir (*Pseudotsuga menziesii*) and sugar pine (*Pinus lambertiana*), *Plant Cell Rep*, 4 (1985) 177-179.
- Murashige T & Skoog F, A revised medium for rapid growth and bioassay with tobacco tissue cultures, *Physiol Plant*, 15 (1962) 473-497.
- Webb D T & Flinn B S, Eastern white pine (*Pinus strobus* L.), in *Biotechnology in agriculture and forestry*, vol 16, Trees III edited by Y P S Bajaj (Springer-Verlag, Berlin) 1991, 358-382.
- Finer J, Kriebel J & Becwar M R, Initiation of embryogenic callus and suspension cultures of eastern white pine (*Pinus strobus* L.), *Plant Cell Rep*, 8 (1989) 203-206.
- Krogstrup P, Effect of culture densities on cell proliferation and regeneration from embryogenic cell suspensions of *Picea sitchensis*, *Plant Sci*, 72 (1990) 115-123.
- Skriver K & Mundy J, Gene expression in response to abscisic acid and osmotic stress, *Plant Cell*, 2 (1990) 503-512.
- Gupta P K & Durzan D J, Biotechnology of somatic polyembryogenesis and plantlet regeneration in loblolly pine, *Bio/Technology*, 5 (1987) 147-151.
- Tremblay L & Tremblay F M, Maturation of black spruce somatic embryos; sucrose hydrolysis and resulting osmotic pressure of the medium, *Plant Cell Tissue Organ Cult*, 42 (1995) 39-46.
- Carrier D J, Cunningham J E, Taylor D C & Dunstan D I, Sucrose requirements and lipid utilization during germination of interior spruce (*Picea glauca engelmannii* complex) somatic embryos, *Plant Cell Rep*, 16 (1997) 550-554.